

IOWA PRENATAL CARE BARRIERS PROJECT

DATA FROM 2017 SURVEYS

2017 Respondents from All Hospitals in Iowa: 20,919
 2017 Respondents with Babies Born in **Scott County: 384**

Demographic Indicators	State	County
Mother's Age		
Less than 18 years	01%	02%
18-19 years	03%	03%
20-25 years	25%	19%
26-30 years	36%	42%
31-35 years	26%	26%
More than 35 years	09%	08%
Mother's Ethnicity		
Hispanic	07%	09%
Non-Hispanic	93%	91%
White	90%	83%
Black	06%	12%
Asian or Pacific Islander	03%	04%
American Indian or Native Alaskan	01%	01%
Socioeconomic Indicators		
Mother's Education		
Less than high school	07%	07%
High school	21%	23%
Some college	34%	34%
College graduate (Bachelor's degree)	27%	22%
Graduate school	12%	13%
Household Income		
Less than \$10,000	12%	16%
\$10,000 - \$19,999	09%	10%
\$20,000 - \$29,999	09%	11%
\$30,000 - \$39,999	08%	07%
\$40,000 - \$49,999	07%	05%
\$50,000 or more	54%	51%

Information Received in 12 Months Prior to Becoming Pregnant	State	County
Health Care Provider Talked About:		
Birth control	49%	56%
Timing of pregnancy	26%	30%
Spacing of children	11%	12%
Doctor, Nurse, or Other Health Care Provider Talked About:		
Taking folic acid or a multivitamin	44%	46%
Smoking	29%	35%
Drinking alcohol	25%	31%
Your weight	24%	29%
Regular exercise	27%	34%
How romantic relationships affect your health	10%	13%
Oral Health	23%	28%
Did not see a doctor, nurse, or other health care professional for a check-up in the 12 months prior to becoming pregnant	31%	24%
Behavioral Indicators During Pregnancy		
No Birth Control was Used at Time of Conception	86%	85%
Unintended Pregnancy and Did NOT Use Birth Control	70%	70%
Cigarettes Smoked During Last 3 Months of Pregnancy		
Not smoked in past 2 years	78%	75%
Zero cigarettes per day	53%	56%
1-10 per day	41%	38%
11-20 per day	05%	07%
More than 20 per day	01%	<01%
Health Care Services During Pregnancy		
Visited a Dentist, Dental Clinic, or Received Dental Care	56%	58%
Diagnosed with Depression During Pregnancy	03%	04%
Took Prescription Medication for Depression	48%	40%
Received Counseling for Depression During Pregnancy	30%	38%
HIV/AIDS Testing During Pregnancy		
Yes	41%	45%
No	32%	30%
No, Testing was offered, but did not want to be tested	02%	01%
Don't know	25%	23%

Received Flu Shot During Pregnancy

Yes	65%	48%
No, Received flu shot before became pregnant	07%	07%
No, Did not want flu shot during pregnancy	12%	19%
No, Doctor said to wait until after pregnancy	01%	<01%
No, Do not ever get flu shots	14%	23%
No, Flu shot was not available	02%	01%

Hospital Services**State****County****Hospital Staff Encouraged Breastfeeding During First 24 Hours After Birth of the Baby**

Yes	84%	81%
No	13%	16%
Don't remember	02%	04%

During Pregnancy or Time at the Hospital, a Doctor, Nurse, or Other Healthcare Professional Provided Information or Scheduled a Time to Discuss:

Action steps to decrease frustration when infant cries	45%	33%
How to lay the baby down to sleep	58%	43%
Dangers of shaking an infant	43%	30%
Did not receive any information and have not scheduled a time to discuss	30%	41%

Things that may have happened at the hospital where the birth occurred

I breastfed my baby in the hospital	81%	79%
Hospital staff gave me information about breastfeeding	87%	84%
My baby was placed in skin-to-skin contact within the first hour of life	84%	86%
I breastfed in the first hour after my baby was born	64%	64%
Hospital staff helped me learn how to breastfeed	72%	74%
Hospital staff told me to breastfeed whenever my baby wanted	68%	73%
I am feeding my baby	94%	92%

Technical Note: Because percentages were rounded, the sum of values may not equal 100%.

For more information: Call the University of Northern Iowa Center for Social & Behavioral Research: (319) 273-2105.

Data source: Iowa Barriers to Prenatal Care Project, Iowa Department of Public Health, University of Northern Iowa Center for Behavioral Research: 2018 Data.