

MBQIP Measures

Phase One Measures

- Heart Failure
 - HF-1: Discharge Instructions
 - HF-2: Evaluation of LVS function
 - HF-3: ACEI or ARB for LVSD
- Pneumonia
 - PN-3b: Blood Cultures Performed in the Emergency Department Prior to Initial Antibiotic Received in Hospital
 - PN-6: Initial Antibiotic Selection for CAP in Immunocompetent Patient

Phase 2 Measures

- Outpatient 1-7: Hospital Compare CMS Measure (all sub-measures that apply):
 - OP-1: Median Time to Fibrinolysis in the Emergency Department
 - OP-2: Fibrinolytic Therapy Received Within 30 Minutes of ED Arrival in the Emergency Department
 - OP-3: Median Time to Transfer to another Facility for Acute Coronary Intervention in the Emergency Department
 - OP-4: Aspirin at Arrival in the Emergency Department
 - OP-5: Median Time to ECG in the Emergency Department
 - OP-6: Timing of Antibiotic Prophylaxis (Prophylactic Antibiotic Initiated Within One Hour Prior to Surgical Incision) in Surgery
 - OP-7: Prophylactic Antibiotic Selection for Surgical Patients in Surgery
- Hospital Consumer Assessment of Healthcare Providers and Systems (HCAHPS)

Phase 3 Measures

- Pharmacist CPOE/Verification of Medication Orders Within 24 Hours
 - Numerator: Number of electronically entered medication orders for an inpatient admitted to a CAH (acute or swing-bed), verified by a pharmacist or directly entered by a pharmacist within 24 hours.
 - Denominator: Total number of electronically entered medication orders for inpatients admitted to a CAH (acute or swing-bed) during the reporting period.
 - Inclusion Criteria: Inpatients admitted to acute care bed, swing bed; observation patients
 - Exclusion Criteria: Outpatients; ED patients
- Outpatient Emergency Department Transfer Communication (Seven Elements)
 - Pre-Transfer Communication Information
 - Patient Identification
 - Vital Signs
 - Medication-related Information
 - Practitioner generated information
 - Nurse generated information
 - Procedures and tests