

# Tobacco Prevention Community Partnership Snapshot — FY 2016

## Chickasaw Co. Public Health & Home Care Services



Executive Director, Kathy Babcock  
260 East Prospect Street  
New Hampton 50659



Contact for tobacco prevention:  
Kathy Babcock  
kbabcock@iowatelecom.net  
641-394-4053

### Population and Funding

#### Population served:

Chickasaw: 12,264  
Floyd: 16,077  
Mitchell: 10,779

**FY 2016  
funding total:  
\$33,799**



Iowa Department of Public Health  
Division of Tobacco Use Prevention and Control

Division Director:  
Jerilyn Oshel

321 E. 12th Street  
Des Moines IA 50319  
515-281-6225  
<http://idph.iowa.gov/tupac/>

### Tobacco Related-Disease Burden age-adjusted rate/100,000 population

County	Death rate				Hospitalization rate	
	cancer	heart disease	stroke	COPD/emphysema	COPD/emphysema	asthma
Chickasaw	152.1	176.7	30.5	46.2	128.9	83.7
Floyd	167.5	179.2	28	51.4	208.9	133.9
Mitchell	125.1	185.8	26.5	24.6	111.4	61.9
State of Iowa	167.7	172.9	35.0	46.9	176.8	77

### Adult cigarette smoking prevalence

IA Behavioral Risk Factor Surveillance System  
2006-12

Chickasaw: 15% (8.8% - 23.2%)

Floyd: 16% (10.3% - 24%)

Mitchell: 15% (9.5% - 21.8%)

State of Iowa (2014): 18.5%

### Youth any tobacco use prevalence

Iowa Youth Survey, 2014

Chickasaw: 17.8%

Floyd: 23.8%

Mitchell: 19.8%

State of Iowa : 18.1%



Population counts are based on 2014 estimates from the U.S. Census Bureau. Adult prevalence estimates are self-reported rates and confidence intervals of rates of cigarette smoking in Iowans 18 and older, average annual rate. Youth rates are self-reported tobacco use rates in 11th grade students. Less than half of 11th grade students in these counties participated in the survey: Boone, Butler, Clarke, Dallas, Des Moines, Fremont, Keokuk, Jasper, Lee, Mahaska, Polk, Poweshiek, Washington and Wayne. In many cases, the percentages shown here for counties with low participation rates will reflect closely their actual rates. Any youth tobacco use includes e-cigarette use. \*At least 5% of all county inpatient records are missing. Deaths are from 2010-14 for COPD and stroke and 2012-14 for heart disease and cancer. Hospitalizations are from 2009-11.

