

**Contraindications and Precautions to the Use of Influenza Vaccines
Advisory Committee on Immunization Practices, United States, 2021-22 Influenza Season**

Vaccine type	Contraindications	Precautions
Egg-based IIV4s	<ul style="list-style-type: none"> History of severe allergic reaction (e.g., anaphylaxis) to any component of the vaccine¹ or to a previous dose of any influenza vaccine (i.e., any egg-based IIV, cclIV, RIV, or LAIV)⁵ 	<ul style="list-style-type: none"> Moderate or severe acute illness with or without fever History of Guillain-Barré syndrome within 6 weeks of receipt of influenza vaccine
cclIV4	<ul style="list-style-type: none"> History of severe allergic reaction (e.g., anaphylaxis) to a previous dose of any cclIV or any component of cclIV4⁵ 	<ul style="list-style-type: none"> Moderate or severe acute illness with or without fever History of Guillain-Barré syndrome within 6 weeks of receipt of influenza vaccine History of severe allergic reaction to a previous dose of any other influenza vaccine (i.e., any egg-based IIV, RIV, or LAIV)⁶
RIV4	<ul style="list-style-type: none"> History of severe allergic reaction (e.g., anaphylaxis) to a previous dose of any RIV or any component of RIV4⁵ 	<ul style="list-style-type: none"> Moderate or severe acute illness with or without fever History of Guillain-Barré syndrome within 6 weeks of receipt of influenza vaccine History of severe allergic reaction to a previous dose of any other influenza vaccine (i.e., any egg-based IIV, cclIV, or LAIV)⁶
LAIV4	<ul style="list-style-type: none"> History of severe allergic reaction (e.g., anaphylaxis) to any component of the vaccine¹ or to a previous dose of any influenza vaccine (i.e., any egg-based IIV, cclIV, RIV, or LAIV)⁵ Concomitant aspirin or salicylate-containing therapy in children and adolescents⁵ Children aged 2 through 4 years who have received a diagnosis of asthma or whose parents or caregivers report that a health care provider has told them during the preceding 12 months that their child had wheezing or asthma or whose medical record indicates a wheezing episode has occurred during the preceding 12 months Children and adults who are immunocompromised due to any cause, including but not limited to immunosuppression caused by medications, congenital or acquired immunodeficiency states, HIV infection, anatomic asplenia, or functional asplenia (e.g., due to sickle-cell anemia) Close contacts and caregivers of severely immunosuppressed persons who require a protected environment Pregnancy Persons with active communication between the CSF and the oropharynx, nasopharynx, nose, or ear or any other cranial CSF leak Persons with cochlear implants** Receipt of influenza antiviral medication within the previous 48 hours for oseltamivir and zanamivir, previous 5 days for peramivir, and previous 17 days for baloxavir⁷ 	<ul style="list-style-type: none"> Moderate or severe acute illness with or without fever History of Guillain-Barré syndrome within 6 weeks of receipt of influenza vaccine Asthma in persons aged ≥ 5 years Other underlying medical conditions that might predispose to complications after wild-type influenza infection (e.g., chronic pulmonary, cardiovascular [except isolated hypertension], renal, hepatic, neurologic, hematologic, or metabolic disorders [including diabetes mellitus])

Abbreviations: ACIP = Advisory Committee on Immunization Practices; cclIV = cell culture–based inactivated influenza vaccine (any valency); cclIV4 = cell culture–based inactivated influenza vaccine, quadrivalent; FDA = Food and Drug Administration; IIV = inactivated influenza vaccine (any valency); IIV4 = inactivated influenza vaccine, quadrivalent; LAIV = live attenuated influenza vaccine (any valency); LAIV4 = live attenuated influenza vaccine, quadrivalent; RIV = recombinant influenza vaccine (any valency); RIV4 = recombinant influenza vaccine, quadrivalent.

* Vaccination providers should check FDA-approved prescribing information for 2021–22 influenza vaccines for the most complete and updated information, including (but not limited to) indications, contraindications, warnings, and precautions. Package inserts for U.S.-licensed vaccines are available at <https://www.fda.gov/vaccines-blood-biologics/approved-products/vaccines-licensed-use-united-statesexternal icon>.

† When a contraindication is present, a vaccine should not be administered, consistent with ACIP General Best Practice Guidelines for Immunization (Kroger A, Bahta L, Hunter P. General best practice guidelines for immunization: best practices guidance of the Advisory Committee on Immunization Practices [ACIP]. <https://www.cdc.gov/vaccines/hcp/acip-recs/general-recs/index.html>). In addition to the contraindications based on history of severe allergic reaction to influenza vaccines that are noted in the Table, each individual influenza vaccine is contraindicated for persons who have had a severe allergic reaction (e.g., anaphylaxis) to any component of that vaccine. Vaccine components can be found in package inserts. Although a history of severe allergic reaction (e.g., anaphylaxis) to egg is a labeled contraindication to the use of egg-based IIV4s and LAIV4, ACIP recommends that persons with a history of egg allergy may receive any licensed, recommended influenza vaccine that is otherwise appropriate for their age and health status. Those who report having had reactions to egg involving symptoms other than urticaria (e.g., angioedema or swelling, respiratory distress, lightheadedness, or recurrent emesis) or who required epinephrine or another emergency medical intervention should be vaccinated in an inpatient or outpatient medical setting (including but not necessarily limited to hospitals, clinics, health departments, and physician offices), if a vaccine other than cclIV4 or RIV4 is used. Vaccine administration should be supervised by a health care provider who is able to recognize and manage severe allergic reactions.

§ When a precaution is present, vaccination should generally be deferred but might be indicated if the benefit of protection from the vaccine outweighs the risk for an adverse reaction, consistent with ACIP General Best Practice Guidelines for Immunization (Kroger A, Bahta L, Hunter P. General best practice guidelines for immunization: best practices guidance of the Advisory Committee on Immunization Practices [ACIP]. <https://www.cdc.gov/vaccines/hcp/acip-recs/general-recs/index.html>). Providers can consider using the following vaccines in these instances; however, vaccination should occur in an inpatient or outpatient medical setting with supervision by a health care provider who is able to recognize and manage severe allergic reactions: 1) for persons with a history of severe allergic reaction (e.g., anaphylaxis) to any egg-based IIV or LAIV of any valency, the provider can consider administering cclIV4 or RIV4; 2) for persons with a history of severe allergic reaction (e.g., anaphylaxis) to any cclIV of any valency, the provider can consider administering RIV4; and 3) for persons with a history of severe allergic reaction (e.g., anaphylaxis) to any RIV of any valency, the provider can consider administering cclIV4. Providers can also consider consulting with an allergist to help determine which vaccine component is responsible for the allergic reaction.

Adapted from **Prevention and Control of Seasonal Influenza with Vaccines: Recommendations of the Advisory Committee on Immunization Practices, United States, 2021–2022 Influenza Season**. The full article is available [here](#).