

Iowa Department of Public Health
BUREAU OF EMERGENCY AND TRAUMA SERVICES

Statewide
EMS TREATMENT PROTOCOLS
ADULT & PEDIATRIC

January 2016



**LUCAS STATE OFFICE BUILDING
DES MOINES, IOWA 50319-0075**

(515) 281-0620

(800) 728-3367

<http://idph.iowa.gov/BETS/EMS>

IOWA EMS TREATMENT PROTOCOLS

Table of Contents

Contents

Protocol Authorization.....	2
Protocol Medications.....	3
2016 Protocol Revisions.....	4
Medical Director Statement of Approval.....	4
Iowa EMS Treatment Protocols	5
Adult Treatment Protocols	5
Pediatric Treatment Protocols	47
Appendices.....	77

Protocol Authorization

Iowa Administrative Code 641 - Chapter 132: Emergency Medical Services—Service Program Authorization

132.8(3) Service program operational requirements. Ambulance and nontransport service programs shall:

b. Utilize department protocols as the standard of care. The service program medical director may make changes to the department protocols provided the changes are within the EMS provider's scope of practice and within acceptable medical practice. A copy of the changes shall be filed with the department.

132.9(2) The medical director's duties include, but need not be limited to:

a. Developing, approving and updating protocols to be used by service program personnel that meet or exceed the minimum standard protocols developed by the department.

Purpose

The completed protocol approval page allows for a physician medical director to implement the use of the *2016 Iowa Statewide EMS Treatment Protocols* for one or more service programs where they serve as the program's medical director.

Instructions

Print or type each service's name in the space below that the medical director is approving the use of the 2016 protocols for. Next select each service's corresponding service type and level of authorization. If the medical director makes any additions, subtractions, or other changes to the 2016 protocols the changes will need to be noted in the Protocol Revisions space and filed with the Department. This would include the addition, subtraction, or change of any medication listed within the 2016 protocols. If no changes are made to the 2016 protocols check the box for no changes.

Scope of Practice

The *Iowa Emergency Medical Care Provider Scope of Practice* document outlines the skills each level of certified EMS provider can perform. Some skills will require the approval of the service program's physician medical director as well as documentation of additional training. Iowa EMS providers may not perform skills outside of their identified scope of practice as documented in the *Iowa Emergency Medical Care Provider Scope of Practice*. The most current version of the *Iowa Emergency Medical Care Provider Scope of Practice* document can be viewed and downloaded from the Bureau's website at: <http://idph.iowa.gov/bets/ems/scope-of-practice>.

Recommendations

It is recommended that each service program maintain records that document the review/education of all staff members on the program's most current protocols and the most current version of the *Iowa Emergency Medical Care Provider Scope of Practice* document.

Service(s) Name		Service Name				
Service Type	Ambulance					
	Nontransport					
Service's Level of Authorization	EMR					
	EMT					
	EMT-I					
	AEMT					
	EMT-P					
	Paramedic					

Protocol Medications

Check All Medications Carried by the Service		
<i>Medication kit should contain <u>only</u> medications approved by the service's Medical Director</i>		
OTC Medications	Medications	
<input type="checkbox"/> Aspirin	<input type="checkbox"/> Adenosine	<input type="checkbox"/> Lorazepam
<input type="checkbox"/> Activated Charcoal	<input type="checkbox"/> Albuterol	<input type="checkbox"/> Magnesium Sulfate
<input type="checkbox"/> Glucose Paste	<input type="checkbox"/> Amiodarone	<input type="checkbox"/> Midazolam
Patient Assisted Medications	<input type="checkbox"/> Atropine	<input type="checkbox"/> Morphine Sulfate
<input type="checkbox"/> Auto-injector Epinephrine	<input type="checkbox"/> Dextrose	<input type="checkbox"/> Naloxone
<input type="checkbox"/> Nitroglycerin	<input type="checkbox"/> Diazepam	<input type="checkbox"/> Nitroglycerin
<input type="checkbox"/> Inhaler	<input type="checkbox"/> Diphenhydramine	<input type="checkbox"/> Ondansetron
IV Fluids	<input type="checkbox"/> Dopamine	<input type="checkbox"/> Oxygen
<input type="checkbox"/> Normal Saline	<input type="checkbox"/> Epinephrine	<input type="checkbox"/> Procainamide
<input type="checkbox"/> Ringer's Lactate	<input type="checkbox"/> Fentanyl	<input type="checkbox"/> Sodium Bicarbonate
<input type="checkbox"/> 5% Dextrose	<input type="checkbox"/> Glucagon	<input type="checkbox"/> Thiamin
	<input type="checkbox"/> Lidocaine	<input type="checkbox"/> Vasopressin
Medications Added by Service's Medical Director		

2016 Protocol Revisions

No changes were made to the *2016 Iowa Statewide EMS Treatment Protocols*

OR

List below or attach copies of all changes made by the physician medical director to the *2016 Iowa Statewide EMS Treatment Protocols*

Page	Protocol Name	Changes Made

Additional Skills for the EMR, EMT, AEMT, & EMT-I

Approval of these additional skills must be within the Service Program's Level of Authorization and the Iowa EMS Provider's Scope of Practice	Mark "Yes" if the skill is approved by the medical director to be performed by the identified certification level	Certification Level	Yes	No
	Pulse oximetry	EMR		
	Glucose monitor	EMT, EMT-I		
	Service carries auto-inject epi	EMT, EMT-I		
	Central line access	EMT-I, AEMT		
	CPAP	EMT, AEMT		

NOTE: Iowa's Scope of Practice document requires medical director approval and documentation of additional training for these skills. Service program must maintain documentation of the additional training

Medical Director Statement of Approval

As the physician medical director I have reviewed both the *2016 Iowa Statewide EMS Treatment Protocols* and the *Iowa Emergency Medical Care Provider Scope of Practice* document and approve the use of the skills, medications, and protocols with revisions as documented above for the authorized Iowa EMS program(s) listed within this document.

Medical Director's Printed Name	Signature	Date

Iowa EMS Treatment Protocols

Section 2

Adult Treatment Protocols

Contents

Initial Patient Care Protocol	6
Abdominal Pain	8
Acute Coronary Syndrome	9
Airway.....	11
Altered Mental Status	12
Amputated Part.....	13
Apparent Death.....	14
Asthma and COPD	15
Behavioral Emergencies	16
Burns.....	17
Cardiac Arrhythmias	20
Childbirth.....	23
Congestive Heart Failure	24
Frostbite	26
Heat Illness	27
Hypothermia.....	28
Nausea & Vomiting.....	29
Pain Control	30
Poisoning	32
Post Resuscitation with Return of Spontaneous Circulation.....	33
Seizure	34
Selective Spinal Immobilization.....	35
Sexual Assault.....	37
Shock	38
Stroke	43
Trauma	44

Initial Patient Care Protocol

REVISED 2016

1. Scene Size Up
 - a) Review the dispatch information
 - b) As you approach the scene consider safety for yourself and your patient.
 - c) Observe universal precautions
 - d) After determining the number and location of patients, consider the need for additional resources
 - e) Determine mechanism of injury and/or nature of illness
 - f) Reassess the situation often

2. Primary Survey
 - a) Obtain general impression of patient, chief complaint, and priority problems
 - b) Determine responsiveness
 - c) Assess airway
 - d) Assess breathing
 - e) Assess circulation

3. Initial Interventions
 - a) Treat airway/breathing problems
 - b) Treat circulation problems
 - c) Establish IV/IO access if indicated
 - d) Apply cardiac monitor if indicated
 - e) Apply pulse oximetry or EtCO₂ monitor if available and indicated
 - f) Treat pain or nausea if present
 - g) Maintain normal patient temperature

4. Secondary Survey
 - a) Perform secondary assessment after initial interventions are completed
 - b) Address problems identified in the secondary survey utilizing the appropriate protocol(s)
 - c) Obtain vital signs., including blood glucose if available and indicated

5. Ongoing Assessment
 - a) Repeated evaluation of patient
 - Vitals every 5 minutes for unstable patients
 - Vitals every 15 minutes for stable patients
 - b) Assess effect of interventions

Initial Patient Care Protocol (continued)

6. Transport/Contact Medical Control
 - a) Patients should be transported as soon as feasible to an appropriate medical facility. Immediate transport with treatment en route is recommended for patients with significant trauma or unstable airways
 - b) Triage with an appropriate service if level of care indicates or assistance is needed and can be accomplished in a timely manner
 - c) Contact medical direction as soon as feasible in accordance with local protocol for further orders
 - d) For seriously injured or critically ill patients, give a brief initial report from the scene when possible, with a more detailed report given to medical direction while en route

Abdominal Pain

(non-traumatic)

REVISED 2012

1. Follow Initial Care Protocol for all Patients

BASIC CARE GUIDELINES

- a) Give nothing by mouth

ADVANCED CARE GUIDELINES

- b) Consider a fluid bolus if indicated.
- c) Evaluate the need for pain and nausea control

Acute Coronary Syndrome

REVISED 2012

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Place patient in position of comfort, loosen tight clothing and provide reassurance. If patient is complaining of shortness of breath, has signs of respiratory distress and pulse oximetry of less than 94% then titrate oxygen to maintain a saturation of 94-99%
- b) If capability exists, obtain a 12-lead EKG and transmit to the receiving facility and/or medical control for interpretation as soon as possible
- c) If patient is alert and oriented and expresses no allergy to aspirin have patient chew nonenteric aspirin 160 – 325 mg
- d) An initial management goal should be to identify STEMI and transport the patient with cardiac symptoms to the facility most appropriate for their needs
- e) Contact medical direction for orders
- f) If the patient has been prescribed nitroglycerin (patient's nitro only) and systolic blood pressure is 90 mmHg or above, give one dose. If patient is taking erectile dysfunction drugs such as Viagra, contact medical direction prior to giving nitroglycerin
- g) Repeat one dose of nitroglycerin in 3-5 minutes if pain continues, systolic blood pressure is 90 mmHg or above and authorized by medical direction, up to a maximum of three doses
- h) If systolic blood pressure less than 90 mmHg or patient does not have prescribed nitroglycerin, transport promptly continuing assessment and supportive measures
- i) Further assess the patient and evaluate the nature of pain (unless other treatment priorities exist). Refer to Appendix F (Reperfusion Strategies) as ordered by medical control.

Acute Coronary Syndrome (Continued)

ADVANCED CARE GUIDELINES

- j) If capability exists, obtain a 12-Lead EKG and transmit to the receiving facility and/or medical control for interpretation as soon as possible
- k) Establish IV access at TKO rate unless otherwise ordered or indicated
- l) Monitor EKG and treat dysrhythmias following appropriate protocols approved by the medical director, referencing AHA guidelines
- m) Administer nitroglycerin (tab or spray) 0.4 mg sublingually if systolic blood pressure 90 mmHg or above for symptoms of chest pain or atypical cardiac pain. Repeat one dose in 5 minutes if pain continues and systolic blood pressure is greater than 90 mmHg or above. Up to a maximum of three doses should be tried before administering morphine sulfate
- n) If pain continues after administration of nitroglycerin and systolic blood pressure remains above 90 mmHg administer morphine sulfate following the AHA ST Elevated Myocardial Infarction (STEMI) guidelines:
 - STEMI – Morphine 2-4 mg IV may repeat 2-8 mg IV every 5 minutes titrated to pain relief and vitals remain stable
 - OR**
 - UA/NSTEMI – Morphine 1-5 mg IV given once

Airway

REVIEWED 2016

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

Breathing spontaneous on initial assessment and adequate ventilation present

- a) Maintain oxygenation with cannula or mask if oxygen saturations are below 94% titrate to 94% - 99%

Breathing spontaneous on initial assessment without adequate ventilation present

- a) Check airway for obstruction and clear if needed
- b) After airway is clear, assist ventilation with an appropriate adjunct and oxygen
- c) If adequate ventilation is not maintained, proceed to an advanced airway

Not breathing on initial assessment

- a) Open airway with head tilt chin lift. If successful, assist ventilations at an adequate rate and depth then reassess
- b) If head tilt chin lift is not successful, check airway for obstruction and clear if needed
- c) After airway is clear, assist ventilation
- d) If adequate ventilation is not maintained, proceed to an advanced airway

Altered Mental Status

REVIEWED 2011

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Obtain blood glucose
- b) If conscious & able to swallow, administer glucose 15 gm by mouth

ADVANCED CARE GUIDELINES

- c) If blood sugar less than 60 mg/dL administer D50 12.5 - 25 gm IV
- d) If no vascular access administer glucagon 1 mg IM
- e) Evaluate the need for naloxone 1 mg IV. If no response may repeat in 3 minutes
- f) Evaluate the need for intubation

Amputated Part

REVISED 2016

1. Follow Initial Patient Care Protocol
2. Follow Trauma Protocol if indicated

BASIC CARE GUIDELINES

- a) Locate amputated part if possible
- b) Wrap amputated part in saline moistened gauze
- c) Place wrapped amputated part in empty plastic bag
- d) Place the plastic bag with the amputated part in a water and ice mixture
- e) Do not use ice alone or dry ice
- f) Label with patient name, the date, and time
- g) Make sure the part is transported with the patient, if possible

Apparent Death

REVISED 2012

1. Follow Initial Patient Care Protocol

Apparent death indications are as follows:

- Signs of trauma are conclusively incompatible with life
- Physical decomposition of the body
- Rigor mortis and/or dependent lividity

If apparent death is confirmed, continue as follows:

BASIC CARE GUIDELINES

- a) The county Medical Examiner and law enforcement shall be contacted
- b) Where possible, contact Iowa Donor Network at 800-831-4131. See Protocol Appendix J
- c) At least one EMS provider should remain at the scene until the appropriate authority is present
- d) Provide psychological support for grieving survivors
- e) Document the reason(s) no resuscitation was initiated
- f) Preserve the crime scene if present
- g) In all other circumstances (except where “NO CPR/DNR” protocol applies; see appendix A) full resuscitation must be initiated

Asthma and COPD

REVISED 2016

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) If patient has a physician prescribed hand-held metered dose inhaler:
 - i. Assist patient in administering a single dose if they have not done so already
 - ii. Reassess patient and assist with second dose if necessary per medical direction
- b) Evaluate the need for CPAP, if available

ADVANCED CARE GUIDELINES

- c) Administer albuterol up to 5.0 mg via nebulizer, repeat as needed
- d) Evaluate the need for epinephrine 1:1,000 concentration 0.3 mg IM.
- e) Evaluate the need for CPAP, if available
- f) Evaluate the need for intubation

Behavioral Emergencies

REVISED 2012

1. Follow Initial Patient Care Protocol
2. If there is evidence of immediate danger, protect yourself and others by summoning law enforcement to help ensure safety

BASIC CARE GUIDELINES

- a) Consider medical or traumatic causes of behavior problems
- b) Keep environment calm

ADVANCED CARE GUIDELINES

- c) For severe anxiety, consider a benzodiazepine such as
 - i. Diazepam 2mg IV every 5 minutes up to 10 mg maximum**OR**
 - ii. Diazepam 5-10mg IM
- d) For excited delirium, consider administering ziprasidone (Geodon) 10-20 mg IM, if available, or similar medication

Burns

REVISED 2016

1. Follow Initial Patient Care Protocol
2. Continually monitor the airway for evidence of obstruction
3. Do not use any type of ointment, lotion, or antiseptic
4. Maintain normal patient temperature
5. Transport according to the Out-of-Hospital Trauma Destination Decision Protocol
(Appendix B)

Thermal Burns

BASIC CARE GUIDELINES

- a) Stop the burning process
- b) Estimate percent of body surface area injured and depth of injury
- c) If wound is less than 10% Body Surface Area, cool burn with Normal Saline
- d) Remove smoldering clothing and jewelry and expose area
- e) Cover the burned area with plastic wrap or a clean dry dressing

ADVANCED CARE GUIDELINES

- f) Establish an IV of LR or NS. For severe burns, consider administration of 500 ml bolus
- g) Contact medical control for further fluid administration
- h) Refer to Pain Control protocol

Burns (continued)

Chemical Burns

BASIC CARE GUIDELINES

- a) Brush off powders prior to flushing. Lint roller may also be used to remove powders prior to flushing
- b) Immediately begin to flush with large amounts of water
- c) Continue flushing the contaminated area when en route to the receiving facility
- d) Do not contaminate uninjured areas while flushing
- e) Attempt to identify contaminant

ADVANCED CARE GUIDELINES

- f) Refer to Pain Control protocol

Toxin in Eye

BASIC CARE GUIDELINES

- a) Flood eye(s) with lukewarm water and have patient blink frequently during irrigation. Use caution to not contaminate other body areas
- b) Attempt to identify contaminant

ADVANCED CARE GUIDELINES

- c) Establish a large bore IV if indicated and infuse as patient condition warrants
- d) Refer to Pain Control protocol

Burns (continued)

Electrical Burns

BASIC CARE GUIDELINES

- a) Treat soft tissue injuries associated with the burn with dry dressing
- b) Treat for shock if indicated

ADVANCED CARE GUIDELINES

- c) Refer to Pain Control protocol

Cardiac Arrhythmias

NEW 2013

1. Follow Initial Care Protocol for all Patients

IF NO PULSE

BASIC CARE GUIDELINES

- a) Perform high quality CPR immediately, apply AED and follow device prompts

ADVANCED CARE GUIDELINES

- b) Perform high quality CPR immediately, apply monitor and check rhythm as soon as possible

VENTRICULAR FIBRILLATION OR VENTRICULAR TACHYCARDIA

- a) Defibrillate at manufacturer's specification, immediately resume CPR for two minutes
- b) Organize therapies such as rhythm and pulse checks, defibrillation, IV/IO access, medication administration and airway management around two minute cycles of CPR
- c) Evaluate for treatable causes
- d) Administer a vasopressor as close to the rhythm check as possible
 - a. Administer epinephrine 1:10,000 concentration 1 mg IV or IO every 3-5 minutes
 - b. Consider administration of vasopressin 40 units IV or IO in place of first or second epinephrine dose
- e) Administer an anti-arrhythmic as close to the rhythm check as possible
 - a. Administer amiodarone 300 mg IV or IO, repeat 150 mg in 5 minutes

TORSADES DE POINTES

- a) Administer magnesium sulfate 1-2 g diluted in 10 ml of d5w IV or IO, delivered over 5-20 minutes

Cardiac Arrhythmias (continued)

ASYSTOLE/PEA

- a) Organize therapies such as rhythm and pulse checks, IV/IO access, medication administration and airway management around two minute cycles of CPR
- b) Evaluate for treatable causes
- c) Administer a vasopressor as close to the rhythm check as possible
 - a. Administer epinephrine 1:10,000 concentration 1 mg IV or IO every 3-5 minutes
 - b. Consider administration of vasopressin 40 units IV or IO in place of first or second epinephrine dose

CARDIAC ARRHYTHMIAS WITH PULSE BASIC CARE GUIDELINES

- a) Follow- Acute Coronary Syndrome protocol

ADVANCED CARE GUIDELINES

BRADYCARDIA

- b) If symptomatic, administer atropine 0.5 mg IV or IO every 3-5 minutes as needed to maximum dose of 3.0 mg
- c) Consider transcutaneous pacing (consider sedation)
OR
Consider administering dopamine 2-10 mcg/kg/min IV or IO
OR
Consider administering epinephrine 2-10 mcg/min IV or IO

TACHYCARDIA (Rates greater than 150)

- a) If patient unstable:
 - a. Perform synchronized cardioversion (consider sedation)
 - b. If narrow complex, consider administration of adenosine 6 mg IV, may be repeated at 12 mg after two minutes
- b) If patient stable
 - a. With wide QRS
 - i. If regular and monomorphic, consider administration of adenosine 6 mg IV, may be repeated at 12 mg after two minutes

- ii. Consider administration of amiodarone 150 mg over 10 minutes
IV or IO
- b. With narrow QRS
 - i. Perform vagal maneuvers
 - ii. Consider administration of adenosine 6 mg IV, may be repeated
at 12 mg after two minutes

Childbirth

REVISED 2012

1. Follow Initial Patient Care Protocol

Normal Delivery

BASIC CARE GUIDELINES

- a) If delivery is imminent with crowning, commit to delivery on site and contact medical control
- b) If the amniotic sac does not break, or has not broken, use a clamp to puncture the sac and push it away from the infant's head and mouth as they appear
- c) For newborn management, see newborn resuscitation protocol

Abnormal Deliveries:

BASIC CARE GUIDELINES

Breech delivery: (buttocks presentation)

- a) Allow spontaneous delivery
- b) Support infant's body as it's delivered
- c) If head delivers spontaneously, proceed as in Section I (Normal Delivery)
- d) If head does not deliver within 3 minutes, insert gloved hand into the vagina, keeping your palm toward baby's face; form a "V" with your fingers and push wall of vagina away from baby's face, thereby creating an airway for baby
- e) Do not remove your hand until relieved by advanced EMS or hospital staff

Congestive Heart Failure

REVISED 2012

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Place patient in position of comfort, typically sitting up, loosen tight clothing and reassure
- b) Maintain oxygenation with cannula or mask if oxygen saturations are below 94% titrate to 94% - 99%
- c) Transport immediately if the patient has any of the following:
 - No history of cardiac problems
 - Systolic blood pressure of less than 100.
 - A history of cardiac problems, but does not have nitroglycerin
- d) If capability exists, obtain a 12-lead EKG and transmit it to the receiving facility and/or medical control for interpretation prior to patient's arrival
- e) Contact medical direction for orders
- f) If the patient has been prescribed nitroglycerin (patient's nitro only) and systolic blood pressure is 90 mmHg or above, give one dose. If patient is taking erectile dysfunction drugs such as Viagra, contact medical direction prior to giving nitroglycerin
- g) Repeat one dose of nitroglycerin in 3-5 minutes if pain continues if systolic blood pressure is 90 mmHg or above and authorized by medical direction, up to a maximum of three doses
- h) Reassess patient and vital signs after each dose of nitroglycerin
- i) Further assess the patient and evaluate possible causes (unless other treatment priorities exist)

ADVANCED CARE GUIDELINES

- j) If not already preformed, obtain a 12-lead EKG and if possible transmit it to the receiving facility and/or medical control
- k) Establish IV access at TKO rate unless otherwise ordered or indicated

Congestive Heart Failure (continued)

- l) Be prepared to intubate patient
- m) Monitor EKG and treat dysrhythmias following the appropriate protocol(s)
- n) Refer to Appendix F (Reperfusion Strategies)
- o) If capability exists, apply CPAP
- p) Administer nitroglycerin (tab or spray) 0.4 mg sublingually if systolic blood pressure 90 mmHg or above

Frostbite

REVIEWED 2011

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Remove the patient from the cold environment
- b) Protect the cold injured extremity from further injury (manual stabilization)
- c) Remove wet or restrictive clothing
- d) Do not rub or massage
- e) Do not re-expose to the cold
- f) Remove jewelry
- g) Cover with dry clothing or dressings

ADVANCED CARE GUIDELINES

- h) Establish IV access at a TKO rate. Use warmed IV fluid if possible
- i) Refer to pain control protocol

Heat Illness

REVIEWED 2011

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Remove from the hot environment and place in a cool environment (back of air conditioned response vehicle)
- b) Loosen or remove clothing
- c) Place in recovery position
- d) Initially cool patient by fanning
- e) Additionally cool patient with cold packs to neck, groin and axilla
- f) If alert, stable and not nauseated, you may have the patient slowly drink small sips of water
- g) If the patient is unresponsive or is vomiting, transport to an appropriate medical facility with patient on their left side

ADVANCED CARE GUIDELINES

- h) Monitor EKG and treat dysrhythmias following the appropriate protocol(s)

Hypothermia

REVIEWED 2011

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Remove wet clothing
- b) If able, check core temperature
- c) Handle patient very gently
- d) Cover patient with blankets

ADVANCED CARE GUIDELINES

- e) Administer warm IV fluids if available, do not administer cold fluids

Nausea & Vomiting

REVIEWED 2011

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Give nothing by mouth

ADVANCED CARE GUIDELINES

- b) Consider fluid bolus IV/IO if evidence of hypovolemia and lung sounds are clear
- c) If patient nauseated or is vomiting, consider anti-emetic medication such as ondansetron (Zofran) 4 mg IV
- d) Consider intubating patients with altered mental status who are vomiting and cannot protect their airway

Pain Control

REVIEWED 2012

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) First, attempt to manage all painful conditions:
 - a) Splint extremity injuries
 - b) Place the patient in a position of comfort

ADVANCED CARE GUIDELINES

- b) Consider administration of pain medications for patients that have significant pain, do not have a decreased level of consciousness, are hemodynamically stable, and have oxygen saturations above 94% medication. Example:
 - a) Morphine 2-4 mg via IV, repeated in 5 min
OR
 - b) Fentanyl 25 to 50 mcg IV every 5 minutes as needed to a maximum of 100 mcg
- c) Administer naloxone 1 mg IV for respiratory depression from narcotics. May repeat once if needed
- d) For severe pain consider anxiolytic medication
 - a) Midazolam 0.5-2.5 mg IV / IM repeated every 5 minutes as needed to a maximum of 5 mg
OR
 - b) Diazepam 2-5 mg IV / IM repeated every 5 minutes as needed to a maximum of 10 mg
OR
 - c) Lorazepam 2mg IV, repeated every 30 minutes as needed to a maximum of 4 mg. Use for long transports
- e) Monitor ECG and O2 saturations

Pain Control (Continued)

- f) The patient must have vital signs taken prior to each dose and be monitored closely. If at any time there is a decreased level of consciousness, decrease in oxygen saturation below 92%, or blood pressure drops to 100 mmHg or less, administration of narcotic medication must stop

Poisoning

REVIEWED 2011

1. Follow Initial Patient Care Protocol
2. Identify contaminate and call Poison Control and follow directions given to provide care: 1-800-222-1222
3. Contact Medical Direction as soon as possible with information given by Poison Control and care given

BASIC CARE GUIDELINES

Ingested poisons

- a) Identify and estimate amount of substance ingested

Inhaled poisons:

- a) Remove patient to fresh air
- b) Administer high flow oxygen.
- c) Estimate duration of exposure to inhaled poison

Absorbed poisons

- a) Identify contaminate! If it will be a hazard to you, use protective clothing and extreme caution

Injected poisons

- a) Be alert for respiratory difficulty. Maintain airway and give high flow oxygen
- b) Check patient for marks, rashes, or welts
- c) Try to identify source of injected poison

Post Resuscitation with Return of Spontaneous Circulation

NEW 2013

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Maintain oxygen saturation between 94% - 99%

ADVANCED CARE GUIDELINES

- b) Consider advanced airway
- c) If available, perform waveform capnography, maintaining PETCO₂ 35-40 mm Hg
- d) Treat hypotension (SBP <90 mm Hg)
 - i. Administer 1-2 L NS or LR
 - ii. Administer dopamine 5-10 mcg/kg/min
 - iii. If bradycardic, consider administration of epinephrine 0.1-0.5 mcg/kg/min
- e) If unable to follow commands, consider induced hypothermia if it can be maintained by receiving hospital
- f) If available, obtain 12-lead EKG
- g) If available, obtain blood glucose
 - i. If blood sugar less than 60 mg/dL administer D50 12.5 - 25 grams IV or IO

Seizure

REVIEWED 2012

1. Follow Initial Patient Care Protocol

Active seizure

BASIC CARE GUIDELINES

- a) Protect airway
- b) Check blood glucose level, if available, and treat hypoglycemia if present

ADVANCED CARE GUIDELINES

- c) Administer valium titrate 2 mg IV push until seizure stops or maximum dose of 10 mg is given
OR
Administer lorazepam 1 mg IV push, titrating 1 mg at a time until the seizure stops or until maximum dose of 10 mg is given

Post seizure

BASIC CARE GUIDELINES

- a) Protect airway
- b) Check blood sugar and treat hypoglycemia if present

ADVANCED CARE GUIDELINES

- c) Consider thiamine 100 mg IM

Selective Spinal Immobilization

Revised 2015

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

2. Patient Presentation:

- a) This protocol is intended for patients who present with a traumatic mechanism of injury.
- b) Immobilization is contraindicated for patients who have penetrating trauma who do not have a neurological deficit.

3. Patient Management:

- a) Assessment:
 - i. Assess for mental status, neurological deficits, spinal pain, tenderness, any evidence of intoxication, or other severe injuries.
 - ii. While maintaining spinal alignment, examine the spine for tenderness on palpation or deformities.
- b) Treatment and Interventions:
 - i. Apply cervical restriction if there is any of the following:
 1. Patient complains of neck pain.
 2. Any neck tenderness of palpation.
 3. Any abnormal mental status, including extreme agitation, or neurological deficit.
 4. Any evidence of alcohol or drug intoxication
 5. There are other severe or painful injuries present.
 6. Any communication barrier that prevents accurate assessment.

Selective Spinal Immobilization (continued)

- ii. Immobilize Patient with cervical collar and a long spine board, full body vacuum splint, scoop stretcher, or similar device if:
 - 1. Patient complains of midline back pain
 - 2. Any midline back tenderness

Note 1: Distracting injuries or altered mental status does not necessitate long spine board use.

Note 2: Patients should not routinely be transported on long boards, unless the clinical situation warrants long board use. An example of this may be facilitation of multiple extremity injuries or an unstable patient where removal of a board will delay transport and/or other treatment priorities. In these rare situation, long boards should be padded or have a vacuum mattress applied to minimize secondary injury to the patient.

Sexual Assault

REVIEWED 2011

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Identify yourself to the patient, assure patient that they are safe and in no further danger
- b) Do not burden patient with questions about the details of the crime; you are there to provide emergency medical care
- c) Be alert to immediate scene and document what you see. Touch only what you need to touch at the scene
- d) Do not disturb any evidence unless necessary for treatment of patient. (If necessary to disturb evidence, document why and how it was disturbed.)
- e) Preserve evidence; such as clothing you may have had to remove for treatment, and make sure that it is never left unattended at any time, to preserve "chain of evidence"
- f) Contact local law enforcement if not present
- g) Treat other injuries as indicated
- h) Treat for shock if indicated

Shock

REVISED 2016

1. Follow Initial Care Protocol for all Patients
2. Maintain oxygen saturation between 94% - 99%

Hypovolemic External Bleeding

BASIC CARE GUIDELINES

- a) Avoid further heat loss
- b) Splint extremities as needed
- c) Follow Hemorrhage Control Protocol
 - i. Control bleeding with direct pressure. Large gaping wounds may need application of a bulky sterile gauze dressing and direct pressure by hand
 - ii. Consider application of tourniquet if unable to control hemorrhage with direct pressure

ADVANCED CARE GUIDELINES

- d) Establish IV/IO access
- e) If radial pulse is absent or systolic blood pressure is less than 90 mmHg, administer 20ml/kg, up to 250ml, NS or LR. Repeat as needed to until radial pulse returns or systolic blood pressure reaches 90 mmHg.

Hypovolemic Internal Bleeding

BASIC CARE GUIDELINES

- a) Place patient in supine position
- b) Consider use of PASG for lower extremity or pelvis fractures
- c) Consider use of pelvic stabilizer for pelvis fractures

ADVANCED CARE GUIDELINES

- d) Establish IV/IO access
- e) If radial pulse is absent or systolic blood pressure is less than 90 mmHg, administer 20ml/kg, up to 250ml, NS or LR. Repeat as needed to until radial pulse returns or systolic blood pressure reaches 90 mmHg.

Cardiogenic

BASIC CARE GUIDELINES

- a) Place in position of comfort
- b) If capability exists, obtain a 12-lead EKG and transmit it to the receiving facility and/or medical control for interpretation prior to patient's arrival

ADVANCED CARE GUIDELINES

- c) Establish IV/IO access
- d) Obtain 12-lead EKG
- e) Administer dopamine IV or IO at 10-20/mcg/kg/min

Obstructive Shock: Tension Pneumothorax

BASIC CARE GUIDELINES

- a) Place in position of comfort

ADVANCED CARE GUIDELINES

- b) Perform needle decompression

Obstructive Shock: Pericardial Tamponade

BASIC CARE GUIDELINES

- a) Place in position of comfort

Shock (continued)

ADVANCED CARE GUIDELINES

- b) The goal should be to minimize scene time with time critical injuries, including establishing IV access en route.
- c) Administer 20 ml/kg, up to 500ml, NS or LR. Repeat as needed to maintain a systolic pressure of 90 mmHg.

Obstructive Shock: Pulmonary Embolus

BASIC CARE GUIDELINES

- a) Place in position of comfort
- b) Avoid further heat loss

ADVANCED CARE GUIDELINES

- c) Administer 20 ml/kg, up to 500ml, NS or LR. Repeat as needed to maintain a systolic pressure of 90 mmHg
- d) If available, obtain 12-lead EKG
- e) Evaluate the need for pain and nausea control
- f) If patient is alert and oriented and expresses no allergy to aspirin, consider having patient chew nonenteric aspirin 160 – 325 mg
- g) Consider administration of dopamine IV or IO at 10-20/mcg/kg/min if systolic blood pressure is less than 90 mmHg.

Distributive Shock: Neurogenic

BASIC CARE GUIDELINES

- a) Place supine
- b) Avoid further heat loss

Shock (continued)

ADVANCED CARE GUIDELINES

- c) Administer 20 ml/kg, up to 500ml, NS or LR. Repeat as needed to maintain a systolic pressure of 90 mmHg
- d) Consider administering dopamine at 10-20 mcg/kg/min IV or IO
- e) If symptomatic bradycardia is present and does not respond to the treatments above, consider:
 - a. administering atropine 0.5 mg every 5 minutes, up to 3 mg
 - OR
 - b. transcutaneous pacing

Distributive Shock: Anaphylactic

BASIC CARE GUIDELINES

- a) If the patient has a physician prescribed Auto-Inject Epinephrine assist with administering it for signs of anaphylaxis

ADVANCED CARE GUIDELINES

- b) Administer epinephrine 1:1,000 concentration 0.01 mg/kg IM, up to a single dose of 0.5 mg. Maximum total dose 1 mg.
- c) Administer diphenhydramine 25 – 50 mg IV/IM
- d) Administer albuterol 2.5mg by nebulizer if respiratory distress
- e) Evaluate need for early intubation if severe anaphylaxis
- f) For cases of severe anaphylaxis consider administration of epinephrine 1:10,000 concentration 0.3 mg - 0.5 mg IV/IO slowly over 3-5 minutes.

Distributive Shock: Septic Shock

BASIC CARE GUIDELINES

- a) Maintain oxygen saturation between 94% - 99%
- b) Place patient in supine position
- c) If temperature is over 102°F/38.9°C, cool patient (i.e. cool sponges)

Shock (continued)

ADVANCED CARE GUIDELINES

- a) Administer 20 ml/kg, up to 500ml, NS or LR. Repeat as needed to maintain a systolic pressure of 90 mmHg
- b) If temperature is over 102°F/38.9°C, cool patient
- c) Consider administering dopamine at 10-20 mcg/kg/min IV or IO
- d) Consider administering diphenhydramine 25 – 50 mg IV/IM

Stroke

REVISED 2012

3. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Perform a “FAST” Cincinnati Prehospital Stroke Scale - checking facial droop, arm drift, speech, and time of onset. Notify receiving facility as soon as possible if stroke is suspected
- b) If Stroke Screening is positive expedite transport to the hospital
- c) Refer to Appendix G (Reperfusion Strategies)
- d) Check blood glucose, if available

ADVANCED CARE GUIDELINES

- e) If blood sugar less than 60 mg/dL administer D50 12.5 - 25 gm IV
 - i. If no vascular access, administer glucagon 1 mg IM
- f) Monitor patient's level of consciousness and blood pressure every five (5) minutes, and keep patient as calm as possible

Trauma

REVISED 2016

1. Follow Initial Patient Protocol for all patients
2. Follow the Out-of-Hospital Trauma Triage Destination Decision Protocol for the identification of time-critical injuries, method of transport and destination decision for treatment of those injuries
3. The goal should be to minimize scene time with time critical injuries, including establishing IVs en route.

Hemorrhage Control

BASIC CARE GUIDELINES

- a) Control bleeding with direct pressure. Large gaping wounds may need application of a bulky sterile gauze dressing and direct pressure by hand
- b) If direct pressure/pressure dressing is ineffective or impractical, apply a tourniquet to extremity
- c) If bleeding site is not amenable to tourniquet placement (i.e. junctional injury), apply a topical hemostatic agent with direct pressure

ADVANCED CARE GUIDELINES

- a) If radial pulse is absent or systolic blood pressure is less than 90 mmHg, administer 20ml/kg, up to 250ml, NS or LR. Repeat as needed to until radial pulse returns or systolic blood pressure reaches 90 mmHg.

Chest Trauma

BASIC CARE GUIDELINES

- a) Seal open chest wounds immediately. Use occlusive dressing taped down. If the breathing becomes worse, loosen one side of the dressing to release pressure and then reseal
- b) Impaled objects must be left in place and should be stabilized by building up around the object with multiple trauma dressings or other cushioning material
- c) Take care that the penetrating object is not allowed to do further damage

Trauma (continued)

ADVANCED CARE GUIDELINES

- d) If concerned for symptomatic pneumothorax, perform needle decompression.

Abdominal Trauma

BASIC CARE GUIDELINES

- a) Control external bleeding. Dress open wounds to prevent further contamination
- b) Evisceration should be covered with a sterile saline soaked occlusive dressing
- c) Impaled objects should be stabilized with bulky dressings for transport

Head, Neck, and Face Trauma

BASIC CARE GUIDELINES

- a) Place the head in a neutral in-line position unless the patient complains of pain or the head does not easily move into this position
- b) Closely monitor the airway. Provide suctioning of secretions or vomit as needed. Be prepared to log roll the patient if they vomit.
- c) Impaled objects in the cheek may be removed if causing airway problems, or you are having trouble controlling bleeding.
- d) Reassess vitals and Glasgow Coma Score (GCS) frequently
- e) Consider eye shield for any significant eye trauma. If the globe is avulsed, do not put it back into socket; cover with moist saline dressing and then place cup over it.

ADVANCED CARE GUIDELINES

- f) Consider intubation if GCS is less than 8 or airway cannot be maintained
- g) If patient is intubated or has an airway such as Combitube, King, LMA
PETCO₂ levels should be continually monitored and maintained at 33 – 43 mmHg if available

Trauma (continued)

Extremity Injuries

BASIC CARE GUIDELINES

- a) Assess extent of injury including presence or absence of pulse
- b) Establish and maintain manual stabilization of injured extremity by supporting above and below the injury
- c) Remove or cut away clothing and jewelry
- d) Cover open wounds with a sterile dressing
- e) Do not intentionally replace any protruding bones
- f) Apply cold pack to area of pain or swelling
- g) If severe deformity of the distal extremity is cyanotic or lacks pulses, align with gentle traction before splinting, and transport immediately

IOWA EMS TREATMENT PROTOCOLS

Section 3

Pediatric Treatment Protocols

Pediatric Initial Care Protocol.....	48
Pediatric Airway.....	50
Pediatric Allergic Reaction.....	52
Pediatric Altered Mental Status	53
Pediatric Apparent Death.....	54
Pediatric Asthma	55
Pediatric Burns	56
Pediatric Cardiac Arrhythmia	59
Pediatric Nausea & Vomiting	62
Pediatric Near Drowning	63
Newborn Resuscitation & Care	65
Pediatric Pain Control.....	68
Pediatric Poisoning	69
Pediatric Seizure	70
Pediatric Selective Spinal Immobilization.....	71
Pediatric Shock	73
Suspected Child Abuse	74
Pediatric Trauma	75

Pediatric Initial Care Protocol

REVISED 2016

1. Scene Size Up
 - a) Review the dispatch information
 - b) As you approach the scene, be sure to consider safety for yourself and your patient
 - c) Observe universal precautions
 - d) After determining the number and location of patients, consider the need for additional resources
 - e) Determine mechanism of injury and/or nature of illness
 - f) Reassess the situation often

2. Primary Survey
 - a) Obtain general impression of patient, chief complaint, and priority problems
 - b) Determine responsiveness
 - c) Assess airway
 - d) Assess breathing
 - e) Assess circulation
 - f) Maintain cervical stabilization/immobilization if indicated
 - g) Utilize length/weight based tape to determine appropriate medications and equipment

3. Initial Interventions
 - a) Treat airway/breathing problems
 - b) Treat circulation problems
 - c) Establish IV/IO access if indicated
 - d) Treat pain or nausea
 - e) Apply cardiac monitor
 - f) Maintain normal patient temperature

4. Secondary Survey
 - a) Perform secondary assessment after initial interventions are completed
 - b) Address problems identified in the secondary survey utilizing the appropriate protocol(s)
 - c) Obtain vital signs, including blood glucose if available and indicated
 - d) Assess pain

5. Ongoing Assessment
 - a) Repeated evaluation of patient
 - Vitals every 5 minutes for unstable patient
 - Vitals every 15 minutes for stable patients
 - b) Assess effect of interventions

Pediatric Initial Care Protocol (continued)

6. Transport/Contact Medical Control
 - a) Patients should be transported as soon as feasible to an appropriate medical facility. Immediate transport with treatment enroute is recommended for patients with significant trauma or unstable airways
 - b) Tier with an appropriate service if level of care indicates or assistance is needed and can be accomplished in a timely manner
 - c) Contact medical direction as soon as feasible in accordance with local protocol for further orders
 - d) For seriously injured or critically ill patients, give a brief initial report from the scene when possible, with a more detailed report given to medical direction while en route

Pediatric Airway

REVIEWED 2016

1. Follow Initial Patient Care Protocol

Breathing spontaneous on initial assessment with adequate ventilation

BASIC CARE GUIDELINES

- a) Maintain oxygenation with cannula, mask, or blow-by if oxygen saturations are below 94%, titrate to 94% - 99%

Breathing without adequate ventilation or not breathing

BASIC CARE GUIDELINES

- a) Open the airway
- b) Attempt assisted ventilation using an appropriate adjunct with high-flow 100% oxygen. If unable to ventilate, first reposition airway and attempt to ventilate again
- c) If ventilation still unsuccessful, check airway for obstruction and attempt to dislodge with age appropriate techniques

ADVANCED CARE GUIDELINES

- d) If unsuccessful establish direct view of object and attempt to remove it with Magill forceps

If obstruction cleared

BASIC CARE GUIDELINES

- a) Assist ventilation and provide oxygen

ADVANCED CARE GUIDELINES

- b) If adequate ventilation is NOT maintained proceed to an advanced airway as appropriate for patient size

Pediatric Airway (continued)

If obstruction not cleared

ADVANCED CARE GUIDELINES

- a) Attempt endotracheal intubation and try to ventilate the patient
- b) If endotracheal intubation is not successful, perform needle cricothyrotomy and needle insufflation

Pediatric Allergic Reaction

REVISED 2012

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Assess airway via Airway Protocol
- b) If the patient has a physician prescribed auto-injectable epinephrine assist with administration and monitor for signs of anaphylaxis

ADVANCED CARE GUIDELINES

- c) Administer epinephrine 1:1,000 concentration 0.01 mg/kg IM, up to a maximum dose of 0.3 - 0.5 mg
- d) Establish IV access
- e) Administer diphenhydramine at 1.0 mg/kg IV or deep IM, up to a maximum dose of 50 mg
- f) Administer epinephrine 1:10,000 concentration 0.01 mg/kg IV for profound shock , up to a maximum dose of 0.3 - 0.5 mg
- g) Administer albuterol 2.5 mg by nebulizer if in respiratory distress

Pediatric Altered Mental Status

REVISED 2012

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Follow Airway Protocol to ensure adequate ventilation
- b) Obtain blood glucose
- c) Patient conscious- give oral Glucose for children over 2 years of age.

ADVANCED CARE GUIDELINES

- d) Establish IV / IO access

If Hypoglycemic

- e) Patient unconscious; give Dextrose 0.5-1.0 g/kg slowly IV up to 25 grams
- f) Patient unconscious and no IV access; administer Glucagon 0.025 mg/kg IM up to 1 mg maximum
- g) Monitor cardiac rhythm
- h) If no improvement in level of consciousness after glucose administration give naloxone 0.1 mg/kg IV up to maximum dose of 2.0 mg per dose
- i) If there is evidence of shock or a history of dehydration, administer a fluid bolus of normal saline at 20 ml/kg over 10-15 minutes
- j) Reassess patient, if signs of shock persist, bolus may be repeated at the same dose up to two times for a maximum total of 60 ml/kg

Pediatric Apparent Death

REVISED 2012

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

Apparent death indications are as follows:

- Signs of trauma are conclusively incompatible with life
- Physical decomposition of the body
- Rigor mortis and/or dependent lividity

If apparent death is confirmed, continue as follows:

- a) The county Medical Examiner and law enforcement shall be contacted
- b) Where possible contact Iowa Donor Network at 800-831-4131

See protocol appendix J

- c) At least one EMS provider should remain at the scene until the appropriate authority is present
- d) Provide psychological support for grieving survivors
- e) Document reason no resuscitation was initiated
- f) Preserve the crime scene if present
- g) In all other circumstances (except where “NO CPR/DNR” protocol applies) full resuscitation must be initiated

Pediatric Asthma

REVISED 2016

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Use Airway Protocol to evaluate the airway and adequacy of ventilation
- b) If patient has a physician prescribed, hand-held metered dose inhaler, assist with administration
- c) Reassess patient and repeat second dose if necessary per medical direction

ADVANCED CARE GUIDELINES

- d) Administer albuterol 2.5.mg via Nebulizer
- e) Evaluate the need for epinephrine 1:1,000 concentration according to length/weight based tape. Dosage may be repeated once after 5 minutes.
- f) Evaluate the need for intubation.

Pediatric Burns

REVISED 2016

1. Follow Initial Patient Care Protocol
2. Continually monitor the airway for evidence of obstruction
3. Do not use any type of ointment, lotion, or antiseptic
4. Maintain normal patient temperature
5. Transport according to the Out-of-Hospital Trauma Destination Decision Protocol
(Appendix B)

Thermal burns

BASIC CARE GUIDELINES

- a) Stop the burning process
- b) Remove smoldering clothing and jewelry
- c) Prevent further contamination of wounds
- d) Cover the burned area with a clean, dry dressing or plastic wrap
- e) Estimate percent of body surface area injured and estimate the depth of burn as superficial, partial thickness or full thickness

ADVANCED CARE GUIDELINES

- f) Establish an IV of LR or NS. For severe burns, consider administration of 20 ml/kg, not to exceed 500 ml.
- g) Contact medical control for further fluid administration
- h) Treat pain per pain protocol

Pediatric Burns (continued)

Chemical burns

BASIC CARE GUIDELINES

- a) Brush off powders prior to flushing. Lint roller may also be used to remove powders prior to flushing
- b) Immediately begin to flush with large amounts of water. Continue flushing the contaminated area when en route to the receiving facility
- c) Do not contaminate uninjured areas while flushing
- d) Attempt to identify contaminant
- e) Transport according to the Out-of-Hospital Destination Decision Protocol (Appendix B)

ADVANCED CARE GUIDELINES

- f) Treat pain per pain control protocol

Toxin in eye

BASIC CARE GUIDELINES

- a) Flood eye(s) with lukewarm water and have patient blink frequently during irrigation. Use caution to not contaminate other body areas
- b) Continue irrigation until advanced personnel take over
- c) Attempt to identify contaminant
- d) Transport to the most appropriate medical facility

ADVANCED CARE GUIDELINES

- e) Treat pain per pain control protocol

Pediatric Burns (continued)

Electrical burns

BASIC CARE GUIDELINES

- a) Treat soft tissue injuries associated with the burn with dry dressing
- b) Treat for shock if indicated
- c) Transport according to the Out-of-Hospital Destination Decision Protocol (Appendix B)
- d) Estimate percent of body surface area injured and estimate the depth of burn as superficial, partial thickness or full thickness

ADVANCED CARE GUIDELINES

- e) Treat pain per pain control protocol

Pediatric Cardiac Arrhythmia

NEW 2013

1. Follow Initial Care Protocol for Pediatric Patients

IF NO PULSE

BASIC CARE GUIDELINES

- a) Perform high quality CPR immediately, apply AED and follow device prompts

ADVANCED CARE GUIDELINES

- b) Perform high quality CPR immediately, apply monitor and check rhythm as soon as possible

VENTRICULAR FIBRILLATION OR VENTRICULAR TACHYCARDIA

- a) Defibrillate at 2J/kg, immediately resume CPR for two minutes
- b) Organize therapies such as rhythm and pulse checks, defibrillation, IV/IO access, medication administration and airway management around two minute cycles of CPR
- c) Second defibrillation at 4 J/kg
- d) Subsequent defibrillations increasing by 2 J/kg, to a maximum of 10 J/kg, not to exceed maximum adult dose
- e) Evaluate for treatable causes
- f) Administer epinephrine 1:10,000 according to length/weight based tape every 3-5 minutes
- g) Administer amiodarone according to length/weight based tape, may repeat twice

Pediatric Cardiac Arrhythmia

ASYSTOLE/PEA

- a) Organize therapies such as rhythm and pulse checks, IV/IO access, medication administration and airway management around two minute cycles of CPR
- b) Evaluate for treatable causes
- c) Administer epinephrine 1:10,000 according to length/weight based tape every 3-5 minutes as needed

CARDIAC ARRHYTHMIAS WITH PULSE

BASIC CARE GUIDELINES

- a) Maintain oxygenation with cannula or mask if oxygen saturations are below 94% titrate to 94% - 99%
- b) Evaluate for treatable causes

BRADYCARDIA WITH SIGNS OF POOR PERFUSION DESPITE OXYGENATION AND VENTILATION

BASIC CARE GUIDELINES

- a) Start CPR if pulse is less than 60 and altered mental status

ADVANCED CARE GUIDELINES

- b) Administer epinephrine 1:10,000 according to length/weight based tape every 3-5 minutes
- c) Consider administration of atropine according to length/weight based tape

Pediatric Cardiac Arrhythmia

TACHYCARDIA (RATES GREATER THAN 180 IN CHILDREN OR 210 IN INFANTS)

ADVANCED CARE GUIDELINES

- a) If patient unstable:
 - i. Perform synchronized cardioversion according to length based tape
(consider sedation)

- b) If patient stable
 - i. Contact medical control

Pediatric Nausea & Vomiting

REVIEWED 2011

1. Follow Initial Patient Care Protocol

ADVANCED CARE GUIDELINES

- a) Initiate IV access
- b) Consider fluid bolus if evidence of hypovolemia
- c) If patient nauseated or is vomiting administer anti-emetic medication such as ondansetron (Zofran) 0.1 mg/kg IV up to 4 mg maximum
- d) Consider intubating patients with altered mental status who are vomiting and can't protect their airway

Pediatric Near Drowning

REVISED 2012

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Establish patient responsiveness
- b) If cervical spine trauma is suspected, manually stabilize the spine
- c) Assess airway for patency, protective reflexes and the possible need for advanced airway management. Look for signs of airway obstruction
- d) Open the airway using head tilt/chin lift if no spinal trauma is suspected, or modified jaw thrust if spinal trauma is suspected
- e) Suction as necessary
- f) Consider placing an oropharyngeal or nasopharyngeal airway adjunct if the airway cannot be maintained with positioning and the patient is unconscious
- g) Assess breathing. Obtain pulse oximeter reading
- h) If breathing is inadequate, assist ventilation using an appropriate adjunct with high-flow, 100% concentration oxygen
- i) Assess circulation and perfusion
- j) If breathing is adequate, place the child in a position of comfort and maintain oxygenation with cannula, mask or blow-by if oxygen saturations are below 94% titrate to 94% - 99%
- k) Assess mental status
- l) If spinal trauma is suspected, continue manual stabilization, apply a rigid cervical collar, and immobilize the patient on a long backboard or similar device
- m) Expose the child only as necessary to perform further assessments. Maintain the child's body temperature throughout the examination
- n) If the child's condition is stable, perform focused history and detailed physical examination on the scene, then initiate transport

Pediatric Near Drowning (continued)

- o) If the child's condition is stable, perform focused history and detailed physical examination on the scene, then initiate transport

ADVANCED CARE GUIDELINES

- p) If abdominal distention arises, consider placing a gastric tube to decompress the stomach if available
- q) If the airway cannot be maintained by other means, including attempts at assisted ventilation, or if prolonged assisted ventilation is anticipated
- r) Perform sedatives and paralytic agents, to aid with intubation as permitted by medical direction. Confirm placement of endotracheal tube using clinical assessment and end-tidal CO₂ monitoring as per medical direction
- s) Initiate cardiac monitoring and determine rhythm. Consult the appropriate protocol for treatment of specific dysrhythmias. Refer to AHA guidelines
- t) Obtain vascular access. Administer normal saline at a maintenance rate according to weight
- u) If the child's condition is critical or unstable, initiate transport as quickly as possible. Perform focused history and detailed physical examination en route to the hospital if patient status and management of resources permit

Newborn Resuscitation & Care

REVISED 2012

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Suction the airway using a bulb syringe as soon as the head is delivered and before delivery of the body. Suction the mouth first, then the nose
- b) Once the body is fully delivered, dry the baby, replace wet towels with dry ones, and wrap the baby in a thermal blanket or dry towel. Cover the scalp to preserve warmth
- c) Open and position the airway. Suction the airway again using a bulb syringe. Suction the mouth first, then the nose
- d) Assess breathing and adequacy of ventilation
- e) If ventilation is inadequate, stimulate by gently rubbing the back and flicking the soles of the feet
- f) If ventilation is still inadequate after brief stimulation, begin assisted ventilation at 40 to 60 breaths per minute using a bag-valve-mask device with room air. If no improvement after 30-60 seconds, apply 100% oxygen
- g) If ventilation is adequate and the infant displays central cyanosis, administer oxygen at 5 lpm via blow-by. Hold the tubing 1/2 to 1 inch from the nose
- h) If the heart rate is slower than 60 beats per minute after 30 seconds of assisted ventilation with high-flow, 100% concentration oxygen, initiate the following actions:
 - Begin chest compressions at a combined rate of 120/minute (three compressions to each ventilation)

Newborn Resuscitation & Care (continued)

ADVANCED CARE GUIDELINES

- i) If there is no improvement in heart rate after 30 seconds. Perform endotracheal intubation
- j) If there is no improvement in heart rate after intubation and ventilation, administer
 - i. epinephrine 1:1000 concentration at 0.1 mg/kg (maximum individual dose 10.0 mg) via endotracheal tube,
 - ii. or epinephrine 1:10,000 concentration at 0.01 mg/kg (maximum individual dose 1.0 mg) IV/IO
 - iii. Repeat epinephrine at the same dose every 3 to 5 minutes as needed
- k) Initiate transport. Reassess heart rate and respirations en route

If the heart rate is between 60 and 80 beats per minute, initiate the following actions:

- Continue assisted ventilation with high-flow, 100% concentration oxygen. If there is no improvement in heart rate after 30 seconds, initiate management sequence described in step H above, beginning with chest compressions
- Initiate transport. Reassess heart rate and respirations en route

If the heart rate is between 80 and 100 beats per minute, initiate the following actions:

- Continue assisted ventilation with high-flow, 100% concentration oxygen. Stimulate as previously described
- Initiate transport. Reassess heart rate after 15 to 30 seconds

If the heart rate is faster than 100 beats per minute, initiate the following actions:

- Assess skin color. If central cyanosis is still present, continue blow by oxygen. Initiate transport. Reassess heart rate and respirations en route

Newborn Resuscitation & Care (continued)

If thick meconium is present

- Initiate endotracheal intubation before the infant takes a first breath. Suction the airway using an appropriate suction adapter while withdrawing the endotracheal tube. Repeat this procedure until the endotracheal tube is clear of meconium. If the infant's heart rate slows, discontinue suctioning immediately and provide ventilation until the infant recovers

Note: If the infant is already breathing or crying, this step may be omitted

Pediatric Pain Control

REVIEWED 2011

1. Follow Initial Patient Care Protocol
2. First attempt to manage all painful conditions with basic care

BASIC CARE GUIDELINES

- a) Splint extremity injuries
- b) Place the patient in a position of comfort

ADVANCED CARE GUIDELINES

- c) Consider administration of pain medications for patients that have significant pain, do not have a decreased level of consciousness, are hemodynamically stable, and have oxygen saturations above 94% medication

Examples:

- Morphine 0.1 mg/kg (maximum individual dose 10 mg) via intravenous or subcutaneous route

OR

- Fentanyl 1.0 mcg/kg (maximum individual dose 100 mcg) via intravenous route

- d) Monitor ECG and O2 saturations
- e) The patient must have vital signs taken prior to each dose and be monitored closely. Administration of narcotic medication must stop if at any time there is a
 - decreased level of consciousness,
 - decrease in oxygen saturation below 92%
 - blood pressure drops to 100 mmHg or less

After drug administration, reassess the patient using the appropriate pain scale

Pediatric Poisoning

REVIEWED 2011

1. Follow Initial Patient Care Protocol
2. Identify contaminate and call Poison Control and follow directions given to provide care: 1-800-222-1222
3. Contact Medical Direction as soon as possible with information given by Poison Control and care given

BASIC CARE GUIDELINES

Ingested Poisons

- a) Identify and estimate amount of substance ingested

Inhaled Poisons:

- a) Remove patient to fresh air
- b) Administer high flow oxygen
- c) Estimate duration of exposure to inhaled poison

Absorbed Poisons

- a) If it will be a hazard to you, use protective clothing and extreme caution

Injected Poisons

- a) Be alert for respiratory difficulty. Maintain airway and give high flow oxygen
- b) Check patient for marks, rashes, or welts

Pediatric Seizure

REVIEWED 2014

1. Follow Initial Patient Care Protocol

Active Seizure

BASIC CARE GUIDELINES

- a) Assess airway via Airway Protocol
- b) Check blood glucose, if available

ADVANCED CARE GUIDELINES

- c) Establish IV access
- d) Administer Benzodiazepine, dosage according to length/weight based device, to stop seizure. May repeat dose in 5 minutes if still seizing
- e) If blood glucose less than 60 mg/dL give glucose IV/IO, or glucagon IM if no IV access, dosage according to length/weight based device.

Post Seizure

BASIC CARE GUIDELINES

- a) Protect airway
- b) Check blood glucose, if available

ADVANCED CARE GUIDELINES

- c) Establish IV
- d) If blood glucose less than 60 mg/dL give glucose IV/IO, dosage according to length/weight based device.

Pediatric Selective Spinal Immobilization

REVISED 2015

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

2. Patient Presentation:

- a) This protocol is intended for patients who present with a traumatic mechanism of injury.
- b) Immobilization is contraindicated for patients who have penetrating trauma who do not have a neurological deficit.

4. Patient Management:

- a) Assessment:
 - i. Assess for mental status, neurological deficits, spinal pain, tenderness, any evidence of intoxication, or other severe injuries.
 - ii. While maintaining spinal alignment, examine the spine for tenderness on palpation or deformities.
- b) Treatment and Interventions:
 - i. Apply cervical restriction if there is any of the following:
 1. Patient complains of neck pain.
 2. Any neck tenderness of palpation.
 3. Any abnormal mental status, including extreme agitation, or neurological deficit.
 4. Any evidence of alcohol or drug intoxication
 5. There are other severe or painful injuries present.
 6. Any communication barrier that prevents accurate assessment.

Pediatric Selective Spinal Immobilization (continued)

- ii. Immobilize Patient with cervical collar and a long spine board, full body vacuum splint, scoop stretcher, or similar device if:
 1. Patient complains of midline back pain
 2. Any midline back tenderness

Note 1: Distracting injuries or altered mental status does not necessitate long spine board use.

Note 2: Patients should not routinely be transported on long boards, unless the clinical situation warrants long board use. An example of this may be facilitation of multiple extremity injuries or an unstable patient where removal of a board will delay transport and/or other treatment priorities. In these rare situation, long boards should be padded or have a vacuum mattress applied to minimize secondary injury to the patient.

Pediatric Shock

REVISED 2012

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Assess airway via Airway Protocol
- b) Assess circulation and perfusion
- c) Control external bleeding
- d) Assess mental status
- e) Expose the child only as necessary to perform further assessments. Maintain the child's body temperature throughout the examination
- f) Initiate transport. Perform focused history and detailed physical examination en route to the hospital if patient status and management of resources permit

ADVANCED CARE GUIDELINES

- g) Initiate cardiac monitoring
- h) Establish IV access using an age-appropriate large-bore catheter with large-caliber tubing. If intravenous access cannot be obtained in a child younger than six years, proceed with intraosseous access. Do not delay transport to obtain vascular access
- i) Administer a fluid bolus of normal saline at 20 ml/kg over 10 to 15 minutes. Reassess patient after bolus. If signs of shock persist, bolus may be repeated at the same dose up to two times for a maximum total of 60 ml/kg

Suspected Child Abuse

REVIEWED 2011

1. Follow Initial Patient Care Protocol

BASIC CARE GUIDELINES

- a) Approach child slowly to establish rapport (except in life-threatening situations), then perform exam
- b) Treat obvious injuries according to appropriate protocol
- c) Genital exam only if indicated in the presence of blood, known or obvious injury and or trauma
- d) Interview parents separate from child, if possible
- e) Transport if permitted by parents
- f) If parents do not allow transport, notify law enforcement for assistance
- g) Communicate vital information only – additional info can be given to attending RN and/or Physician on arrival
- h) Record observations and factual information on run report
- i) Report all suspected abuse to the National hotline at 1-800-362-2178 within 24 hours of your contact of the patient. This will be an oral report only
- j) Within 48 hours of oral reporting, you must submit a written report for all suspected abuse to the Department of Human Services

Pediatric Trauma

REVISED 2016

1. Follow Initial Patient Care Protocol
2. Follow the Out-of-Hospital Trauma Triage Destination Decision Protocol for the identification of time critical injuries, method of transport and trauma facility resources necessary for treatment of those injuries
3. The goal should be to minimize scene time with time critical injuries, including establishing IVs en route.

BASIC CARE GUIDELINES

- a) Follow Shock Protocol if shock is present

Hemorrhage Control:

BASIC CARE GUIDELINES

- a) Control bleeding with direct pressure. Large gaping wounds may need application of a bulky sterile gauze dressing and direct pressure by hand
- b) If direct pressure/pressure dressing is ineffective or impractical, apply a tourniquet to extremity
- c) If bleeding site is not amenable to tourniquet placement (i.e. junctional injury), apply a topical hemostatic agent with direct pressure

ADVANCED CARE GUIDELINES

- d) Establish large bore IV
- e) Cardiac monitor

Pediatric Trauma (continued)

Chest Trauma:

BASIC CARE GUIDELINES

- a) Seal open chest wounds immediately. Use occlusive dressing taped down. If the breathing becomes worse, loosen one side of the dressing to release pressure and then reseal
- b) Impaled objects must be left in place and should be stabilized by building up around the object with multiple trauma dressings or other cushioning material
- c) Take care that the penetrating object is not allowed to do further damage

Abdominal Trauma

BASIC CARE GUIDELINES

- a) Control external bleeding. Dress open wounds to prevent further contamination
- b) Evisceration should be covered with a sterile saline soaked occlusive dressing
- c) Impaled objects should be stabilized with bulky dressings for transport

Head, Neck, and Face Trauma

BASIC CARE GUIDELINES

- a) Place the head in a neutral in-line position unless the patient complains of pain or the head does not easily move into this position
- b) Closely monitor the airway. Provide suctioning of secretions or vomit as needed. Be prepared to log roll the patient if they vomit. Maintain manual spinal stabilization if patient is log rolled
- c) Reassess vitals, GCS and pupillary response frequently
- d) Consider eye shield for any significant eye trauma. If the globe is avulsed, do not put it back into socket; cover with moist saline dressing and then place cup over it

IOWA EMS TREATMENT PROTOCOLS

Section 4

Appendices

EMS Out-Of-Hospital Do-Not-Resuscitate Protocol	78
Adult Out Of Hospital Trauma Triage Destination Decision Protocol	79
Pediatric Out Of Hospital Trauma Triage Destination Decision Protocol.....	80
Physician On Scene.....	81
Air Medical Transport.....	82
Discontinuation Of Resuscitation	83
Reperfusion Therapy Screening Not Limited To Paramedic Level	84
Strategies for Reperfusion Therapy: Acute Stroke	86
S T A R T	88
Pediatric JumpSTART	89
Guidelines for EMS Provider Initiating Organ & Tissue Donation	90
Guidelines for EMS Providers responding to a patient with special needs	91
EMS Approved Abbreviations.....	92
Guidelines For New Protocol Development	93

Appendix A

EMS Out-of-Hospital Do-Not-Resuscitate Protocol

Purpose: This protocol is intended to avoid unwarranted resuscitation by emergency care providers in the out-of-hospital setting for a qualified patient.¹ There must be a valid Out-Of-Hospital Do-Not-Resuscitate (OOH DNR) order signed by the qualified patient's attending physician or the presence of the OOH DNR identifier indicating the existence of a valid OOH DNR order.

No resuscitation: Means withholding any medical intervention that utilizes mechanical or artificial means to sustain, restore, or supplant a spontaneous vital function, including but not limited to:

1. Chest compressions,
2. Defibrillation,
3. Esophageal/tracheal/double-lumen airway; endotracheal intubation, or
4. Emergency drugs to alter cardiac or respiratory function or otherwise sustain life.

Patient criteria: The following patients are recognized as qualified patients to receive no resuscitation:

1. The presence of the uniform OOH DNR order or uniform OOH DNR identifier, or
2. The presence of the attending physician to provide direct verbal orders for care of the patient.

The presence of a signed physician order on a form other than the uniform OOH DNR order form approved by the department may be honored if approved by the service program EMS medical director. However, the immunities provided by law apply only in the presence of the uniform OOH DNR order or uniform OOH DNR identifier. When the uniform OOH DNR order or uniform OOH DNR identifier is not present contact must be made with on-line medical control and on-line medical control must concur that no resuscitation is appropriate.

Revocation: An OOH DNR order is deemed revoked at any time that a patient, or an individual authorized to act on the patient's behalf as listed on the OOH DNR order, is able to communicate in any manner the intent that the order be revoked. The personal wishes of family members or other individuals who are not authorized in the order to act on the patient's behalf shall not supersede a valid OOH DNR order.

Comfort Care (♥): When a patient has met the criteria for no resuscitation under the foregoing information, the emergency care provider should continue to provide that care which is intended to make the patient comfortable (a.k.a. ♥ **Comfort Care**). Whether other types of care are indicated will depend upon individual circumstances for which medical control may be contacted by or through the responding ambulance service personnel.

♥ **Comfort Care** may include, but is not limited to:

1. Pain medication.
2. Fluid therapy.
3. Respiratory assistance (oxygen and suctioning).

Qualified Patient means an adult patient determined by an attending physician to be in a terminal condition for which the attending physician has issued an Out of Hospital DNR order in accordance with the law. Iowa Administrative Code 641-142.1 (144A) Definitions.

Appendix B

Revised 2014

Adult Out Of Hospital Trauma Triage Destination Decision Protocol

The following criteria shall be utilized to assist the EMS provider in the identification of time critical injuries, method of transport and trauma care facility resources necessary for treatment of those injuries

Step 1 - Assess for Time Critical Injuries: Level of Consciousness & Vital Signs

- Glasgow Coma Score ≤ 13
- Respiratory rate <10 or >29 breaths per minute, or need for ventilatory support.
- Systolic B/P (mmHg) less than <90 mmHg

If ground transport time to a Resource (Level I) or Regional (Level II) Trauma Care Facility is less than 30 minutes, transport to the nearest Resource (Level I) or Regional (Level II) Trauma Care Facility. If greater than 30 minutes, ground transport time to Resource (Level I) or Regional (Level II) Trauma Care Facility, transport to the nearest appropriate Trauma Care Facility. If time can be saved or level of care needs exist, tier with ground or air ALS service program

If step 1 does not apply, move on to step 2

Step 2 - Assess for Anatomy of an Injury

- All penetrating injuries to head, neck, torso and extremities proximal to elbow or knee
- Chest wall instability or deformity (e.g., flail chest)
- Suspected two or more proximal long-bone fractures
- Suspected pelvic fractures
- Crushed, degloved, mangled, or pulseless extremity
- Open or depressed skull fracture
- Amputation proximal to wrist or ankle
- Paralysis or Parasthesia
- Partial or full thickness burns $> 10\%$ TBSA or involving face/airway

If ground transport time to a Resource (Level I) or Regional (Level II) Trauma Care Facility is less than 30 minutes, transport to the nearest Resource (Level I) or Regional (Level II) Trauma Care Facility. If greater than 30 minutes ground transport time to Resource (Level I) or Regional (Level II) Trauma Care Facility, transport to the nearest appropriate Trauma Care Facility. If time can be saved or level of care needs exist, tier with ground or air ALS service program

If step 2 does not apply, move on to step 3

Step 3 - Consider Mechanism of Injury & High Energy Transfer

- Falls
- Adult: > 20 ft. (one story is equal to 10 feet)
- Auto vs. pedestrian/bicyclist thrown, run over, or with significant (>20 mph) impact
- High-risk auto crash:
 - Interior compartment intrusion, including roof: Motorcycle crash >20 mph,
 - >12 inches occupant site; >18 inches any site
 - Ejection (partial or complete) from automobile
 - Death in same passenger compartment
 - Vehicle telemetry data consistent with high risk of injury

Transport to the nearest appropriate Trauma Care Facility, need not be the highest level trauma care facility.

If step 3 does not apply, move on to step 4

Step 4 - Consider risk factors:

- Older adults
- Risk of injury/death increases after age 55 years
- SBP <110 might represent shock after age 65 years
- EMS provider judgment
- Low impact mechanisms (e.g. ground level falls) might result in severe injury
- ETOH/Drug use
- Pregnancy > 20 weeks
- Anticoagulants and bleeding disorders
 - oPatients with head injury are at high risk for rapid deterioration

Transport to the nearest appropriate Trauma Care Facility, need not be the highest level trauma care facility.

If none of the criteria in the above 4 steps are met, follow local protocol for patient disposition. When in doubt, transport to nearest trauma care facility for evaluation.

For all Transported Trauma Patients:

1. Patient report to include: MOI, Injuries, Vital Signs & GCS, Treatment, Age, Gender and ETA
2. Obtain further orders from medical control as needed.

(April 2013)

Appendix B

Revised 2014

Pediatric Out Of Hospital Trauma Triage Destination Decision Protocol

The following criteria shall be utilized to assist the EMS provider in the identification of time critical injuries, method of transport and trauma care facility resources necessary for treatment of those injuries

Step 1 - Assess for Time Critical Injuries: Level of Consciousness & Vital Signs

Abnormal Responsiveness: abnormal or absent cry or speech. Decreased response to parents or environmental stimuli. Floppy or rigid muscle tone or not moving. Verbal, Pain, or Unresponsive on AVPU scale.

OR

Airway/Breathing Compromise: obstruction to airflow, gurgling, stridor or noisy breathing. Increased/excessive retractions or abdominal muscle use, nasal flaring, stridor, wheezes, grunting, gasping, or gurgling. Decreased/absent respiratory effort or noisy breathing. Respiratory rate outside normal range.

OR

Circulatory Compromise: cyanosis, mottling, paleness/pallor or obvious significant bleeding. Absent or weak peripheral or central pulses; pulse or systolic BP outside normal range. Capillary refill > 2 seconds with other abnormal findings.

Glasgow Coma Score ≤13

If ground transport time to a Resource (Level I) or Regional (Level II) Trauma Care Facility is less than 30 minutes, transport to the nearest Resource (Level I) or Regional (Level II) Trauma Care Facility. If greater than 30 minutes, ground transport time to Resource (Level I) or Regional (Level II) Trauma Care Facility, transport to the nearest appropriate Trauma Care Facility. If time can be saved or level of care needs exist, tier with ground or air ALS service program

If step 1 does not apply, move on to step 2

Step 2 - Assess for Anatomy of an Injury

- All penetrating injuries to head, neck, torso and extremities proximal to elbow or knee
- Chest wall instability or deformity (e.g., flail chest)
- Suspected two or more proximal long-bone fractures
- Suspected pelvic fractures
- Crushed, degloved, mangled, or pulseless extremity
- Open or depressed skull fracture
- Amputation proximal to wrist or ankle
- Paralysis or Parasthesia
- Partial or full thickness burns > 10% TBSA or involving face/airway

If ground transport time to a Resource (Level I) or Regional (Level II) Trauma Care Facility is less than 30 minutes, transport to the nearest Resource (Level I) or Regional (Level II) Trauma Care Facility. If greater than 30 minutes ground transport time to Resource (Level I) or Regional (Level II) Trauma Care Facility, transport to the nearest appropriate Trauma Care Facility. If time can be saved or level of care needs exist, tier with ground or air ALS service program

If step 2 does not apply, move on to step 3

Step 3 - Consider Mechanism of Injury & High Energy Transfer

- Falls >10 feet or two times the height of the child
- High-risk auto crash
 - Interior compartment intrusion, including roof: >12 inches occupant site;
 - >18 inches any site
 - Ejection (partial or complete) from automobile
 - Death in same passenger compartment
 - Vehicle telemetry data consistent with high risk of injury
- Auto vs. pedestrian/bicyclist thrown, run over, or with significant (>20 mph) impact
- Motorcycle crash >20 mph

Transport to the nearest appropriate Trauma Care Facility, need not be the highest level trauma care facility.

If step 3 does not apply, move on to step 4

Step 4 - Consider risk factors:

- Pregnancy > 20 weeks
- Anticoagulants and bleeding disorders
 - Patients with head injury are at high risk for rapid deterioration
- EMS provider Judgment
- ETOH/Drug use

Transport to the nearest **(Any Level)** Trauma Care Facility.

If none of the criteria in the above 4 steps are met, follow local protocol for patient disposition. When in doubt, transport to nearest trauma care facility for evaluation.

For all Transported Trauma Patients:

1. Patient report to include: MOI, Injuries, Vital Signs & GCS, Treatment, Age, Gender and ETA
2. Obtain further orders from medical control as needed.

(April 2013)

Appendix C

Physician On Scene

Your offer of assistance is appreciated. However, this EMS service, under law and in accordance with nationally recognized standards of care in Emergency Medicine, operates under the direct authority of a Physician Medical Director. Our Medical Director and physician designees have already established a physician-patient relationship with this patient. To ensure the best possible patient care, and to prevent inadvertent patient abandonment or interference with an established physician-patient relationship, please comply with our established protocols.

Please review the following if you wish to assume responsibility for this patient:

1. You must be recognized or identify yourself as a qualified physician.
2. You must be able to provide proof of licensure and identify your specialty.
3. If requested, you must speak directly with the on-line medical control physician to verify transfer of responsibility for the patient from that physician to you.
4. EMS personnel, in accordance with state law, can only follow orders that are consistent with the approved protocols.
5. You must accompany this patient to the hospital, unless the on-line medical control physician agrees to re-assume responsibility for this patient prior to transport.

Appendix D

Air Medical Transport Utilization Guidelines for Scene Response

These guidelines have been developed to assist with the decision making for use of air medical transport by the emergency medical services community. The goal is to match the patient's needs to the timely availability of resources in order to improve the care and outcome of the patient from injury or illness.

CLINICAL INDICATORS:

1. Advanced level of care need (skills or medications) exists that could be made available more promptly with an air medical tier versus tiering with ground ALS service, and further delay would likely jeopardize the outcome of the patient
2. Transport time to definitive care hospital can be significantly reduced for a critically ill or injured patient where saving time is in the best interest of the patient
3. Multiple critically ill or injured patients at the scene where the needs exceed the means available
4. EMS Provider 'index of suspicion' based upon mechanism of injury and patient assessment

DIFFICULT ACCESS SITUATIONS:

1. Wilderness or water rescue assistance needed
2. Road conditions impaired due to weather, traffic, or road construction / repair
3. Other locations difficult to access

The local EMS provider must have a good understanding of regional EMS resources and strive to integrate resources to assure that ground and air services cooperate as efficiently and effectively as possible in the best interest of the patient.

Medical directors for ambulance services should assure that EMS providers are aware of their own service's abilities and limitations given the level of care and geographic response area being served. Audits should be conducted on an ongoing basis to assure that utilization of regional resources (ground and air) is appropriate in order to provide the level of care needed on a timely basis.

Appendix E

DISCONTINUATION OF RESUSCITATION

INDICATIONS TO CONSIDER TERMINATION OF RESUSCITATION:

1. Patient is in full arrest with no signs of life present.
2. Patient is considered an adult.
3. Full ACLS has been instituted (Paramedic level) to include rhythm analysis and defibrillation if indicated, advanced airway management, and drugs given per protocol.
4. No return of circulation or shockable rhythm exists.
5. Correctable causes or special resuscitation circumstances have been considered and addressed.

TERMINATION OF RESUSCITATION:

1. Patient meets all five criteria under 'indications' above, or patient is terminally ill/DNR where CPR was started prior to knowledge of resuscitation status.
2. *Physician on-line medical direction* is contacted (while ACLS continues) to discuss any further appropriate actions.
3. ACLS may be discontinued if *physician on-line medical direction* authorizes.

OTHER CONSIDERATIONS:

1. Documentation must reflect that the decision to terminate resuscitation was determined by *physician on-line medical direction*.
2. An EMS/health care provider must attend the deceased until the appropriate authorities arrive.
3. All IVs, tubes, etc. should be left in place until the medical examiner authorizes removal.
4. Implement survivor support plans related to coroner notification, funeral home transfer, leaving the body at the scene, and death notification/grief counseling for survivors.

Physician on-line medical direction includes either of the following:

1. Hospital based physician contact via phone or radio.
2. Patient's primary care physician or on call physician contact via phone or radio.

Special Considerations

Patients with profound hypothermia or drug or toxin overdose may benefit from continued resuscitation.

Reperfusion Therapy Screening Not Limited To Paramedic Level

This form should be completed for patients suffering from Acute Coronary Syndromes. This tool will be used to triage patients to the appropriate receiving facility, and provide a template for passing information on to the receiving facility. Fibrinolytic screening may be done at the EMT-B level; however the decision to bypass a local hospital to transport to a Percutaneous Coronary Intervention (PCI) capable facility is reserved for the PS and Paramedic levels.

1. If available, obtain 12-Lead EKG and transmit to receiving facility
2. EMT level – Transport patient to closest appropriate facility. Contact medical control for decision on completing thrombolytic checklist.
3. PS/Paramedic Level – Evaluate 12-Lead for evidence of STEMI.

If STEMI is present, determine appropriate destination.

- If transport time to a facility capable of providing emergency PCI care is 60 minutes or less, it is recommended that all of these patients be transported directly to the emergency PCI capable facility.
- If transport time to a facility capable of providing emergency PCI care is between 60 - 90 minutes, transport to the PCI capable facility should be considered.
- If transport is initiated to a non-PCI facility:
 1. Complete fibrinolytic therapy checklist on next page.
 2. If a local protocol for fibrinolytic therapy in the field has been established, then proceed with fibrinolytic protocol if:
 - Authorized by voice contact with medical control, and
 - The PS/Paramedic has received training and has the approval of their physician medical director

In all instances those patients requiring immediate hemodynamic or airway stabilization should be transported to the closest appropriate facility.

If STEMI is not present, transport patient to closest appropriate facility.

Note: See Fibrinolytic Checklist on the following page

If directed by medical control, complete fibrinolytic checklist below

Fibrinolytic Checklist

Any **YES** findings will be relayed to medical control. **Absolute Contraindications** preclude the use of fibrinolytics. **Relative Contraindications** require consultation with medical control.

DATE:	PATIENT AGE:	MALE	FEMALE	INCIDENT/RECORD #:	YES	NO
ABSOLUTE CONTRAINDICATIONS						
Any known intracranial hemorrhage?						
Known structural cerebral vascular lesion?						
Ischemic stroke within 3 months EXCEPT acute ischemic stroke within 3 hours?						
Suspected aortic dissection?						
Active bleeding or bleeding diathesis (excluding menses)?						
Significant closed head trauma or facial trauma within 3 months?						
RELATIVE CONTRAINDICATIONS						
History of chronic, severe, poorly controlled hypertension?						
Severe, uncontrolled hypertension on presentation (S >180mmHg or D>110mmHg)						
History of prior ischemic stroke >3 months, dementia, or known intracranial pathology?						
Traumatic or prolonged (>10 min) CPR or major surgery (<3 weeks)						
Non-compressible vascular punctures?						
Pregnancy?						
Active peptic ulcer?						
Current use of anticoagulants?						
EMS Provider Print Name:				Signature:		

Appendix G

Strategies for Reperfusion Therapy: Acute Stroke

(NEW 2012)

Reperfusion Therapy Screening Not Limited to Paramedic Level

This appendix should be used for suspected acute stroke. This tool will be used to triage patients to the appropriate receiving facility, and provide a template for passing information to the receiving facility.

1. Perform a Cincinnati pre-hospital stroke screen (or other reproducible stroke assessment).
2. If assessment is positive for stroke, and onset of symptoms can be established within the past 4.5 hours, then determine the appropriate destination:
 - a. If transport time to a Primary Stroke Center is less than 30 minutes, it is recommended that all of these patients be transported directly to the Primary Stroke Center
 - b. If transport time to a Primary Stroke Center is greater than 30 minutes, then transport to the nearest stroke capable hospital.
3. Consider the use of air transport if it will facilitate the arrival of the acute stroke patient for treatment within 4.5 hours to a Primary Stroke Center or stroke capable hospital.
4. If transport to a Primary Stroke Center or stroke capable hospital cannot be achieved to arrive within 4.5 hours, then transport to the closest appropriate facility.
5. In all instances, those patients requiring immediate hemodynamic or airway stabilization should be transported to the closest appropriate facility.
6. Complete the fibrinolytic checklist on next page.

Primary Stroke Center – hospitals that have been certified by the Joint Commission on Hospital Accreditation or an equivalent agency to meet Brain Attack Coalition and American Stroke Association guidelines for stroke care

Stroke capable hospital – hospitals that have the following:

- rt-PA readily available for administration
- Head CT, laboratory and EKG capabilities 24/7
- Process in place for transporting appropriate patients to a Primary Stroke Center
- Stroke protocol in place that follows American Stroke Association guidelines
- Emergency department coverage by physician, or advanced practitioner

The list of Iowa Hospital Triage Destinations is available on the Iowa Hospital Association web site at:
http://www.ihconline.org/UserDocs/Pages/Iowa_Hospital_Stroke_Triage_System.pdf

If directed by medical control, complete fibrinolytic checklist below

Appendix G

Fibrinolytic Checklist

Any **YES** findings will be relayed to medical control. **Absolute Contraindications** preclude the use of fibrinolytics. **Relative Contraindications** require consultation with medical control.

DATE:	PATIENT AGE:	MALE	FEMALE	INCIDENT/RECORD #:	YES	NO
ABSOLUTE CONTRAINDICATIONS						
Any known intracranial hemorrhage?						
Known structural cerebral vascular lesion?						
Ischemic stroke within 3 months EXCEPT acute ischemic stroke within 3 hours?						
Suspected aortic dissection?						
Active bleeding or bleeding diathesis (excluding menses)?						
Significant closed head trauma or facial trauma within 3 months?						
RELATIVE CONTRAINDICATIONS						
History of chronic, severe, poorly controlled hypertension?						
Severe, uncontrolled hypertension on presentation (S >180mmHg or D>110mmHg)						
History of prior ischemic stroke >3 months, dementia, or known intracranial pathology?						
Traumatic or prolonged (>10 min) CPR or major surgery (<3 weeks)						
Non-compressible vascular punctures?						
Pregnancy?						
Active peptic ulcer?						
Current use of anticoagulants?						
EMS Provider Print Name:				Signature:		

Appendix H


Simple Triage and Rapid Treatment

START

The following are guidelines for initial tactical triage using the START method. START is most useful in initially clearing the disaster zone where there are numerous casualties. **It focuses on respiration rate, perfusion, and mental status and takes under one minute to complete.** Once the patient moves toward a higher level of care (evacuation), a more detailed approach to triage may be needed.


Respirations
Perfusion
Mental Status

Green = Minor/Ambulatory
Yellow = Delayed
Red = Immediate
Black = Deceased/Expectant



Appendix I

Pediatric JumpSTART



©Lou Romig MD, 2002

Appendix J

Guidelines for EMS Provider Initiating Organ & Tissue Donation

At the Scene of the Deceased

1. All appropriate patient care protocols will be enacted to assure patient care is provided according to prevailing standards.
2. If resuscitation efforts are unsuccessful or if upon arrival the patient is deceased and without indications to initiate resuscitation, then on-line medical direction will be contacted to confirm that no further medical care is to be given.
3. As per Iowa Code 142C.7 a medical examiner or a medical examiner's designee, peace officer, fire fighter, or emergency medical care provider may release an individual's information to an organ procurement organization, donor registry, or bank or storage organization to determine if the individual is a donor.
4. As per Iowa Code 142C.7 any information regarding a patient, including the patient's identity, however, constitutes confidential medical information and under any other circumstances is prohibited from disclosure without the written consent of the patient or the patient's legal representative.
5. At least one EMS provider should remain at the scene until the appropriate authority (medical examiner, funeral home, public safety, etc.) is present.
6. Contact IOWA DONOR NETWORK at 800-831-4131

Appendix K

Guidelines for EMS Providers responding to a patient with special needs

(This Protocol is not intended for interfacility transfers.)

These guidelines should be used when an EMS provider, responding to a call, is confronted with a patient using specialized medical equipment that the EMS provider has not been trained to use, and the operation of that equipment is outside of the EMS provider's scope of practice. The EMS provider may treat and transport the patient, as long as the EMS provider doesn't monitor or operate the equipment in any way while providing care.

When providing care to patients with special needs, EMS personnel should provide the level of care necessary, within their level of training and certification. When possible, the EMS provider should consider utilizing a family member or caregiver who has been using this equipment to help with monitoring and operating the special medical equipment if necessary during transport.

Some examples of special medical devices:

- PCA (patient controlled analgesic)
- Chest Tube

Appendix L

EMS APPROVED ABBREVIATIONS

ā	before	ER	emergency room	PAC	premature atrial contraction
ABC	airway, breathing, circulation	ET	endotracheal	PAT	paroxysmal atrial tachycardia
ALS	advanced life support	ETOH	alcohol	PCR	patient care record
AMI	acute myocardial infarction	fib	fibrillation	PE	physical exam, pulmonary edema
amps	ampules	fl	fluid	pedi	pediatric
ASA	aspirin	fx	fracture		PERLpupils equal, reactive to light
AT	atrial tachycardia	GI	gastrointestinal	PJC	premature junctional
AV	atrioventricular	gm	gram	po	by mouth
bicarb	sodium bicarbonate	gr	grain	pr	per rectum
BID	twice a day	gt(t)	drop(s)	prn	whenever necessary, as needed
BLS	basic life support	h,hr	hour	PVC	premature ventricular contraction
BP	blood pressure	hx	history	q	every
BS	blood sugar	ICU	intensive care unit	QID	four times a day
Ā	with	IM	intramuscular	R	respirations
CAD	coronary artery disease	IV	intravenous	R/O	rule out
CC	chief complaint	Kg	kilogram	RN	registered nurse
cc	cubic centimeter	KVO	keep vein open	Rx	treatment
CCU	coronary care unit	L	liter	s	without
CHB	complete heart block	LOC	level of consciousness	SC	subcutaneous
CHF	congestive heart failure	LR	lactated ringers	Sec	second
cm	centimeter	Mgtt	microdrip	SL	sublingual
CNS	central nervous system	MD	medical doctor	SOB	shortness of breath
c/o	complains of	mEq	milliequivalents	SQ	subcutaneous
CO	carbon monoxide	mg	milligram	STAT	immediately
CO ₂	carbon dioxide	MI	myocardial infarction	s/s	sign, symptoms
COPD	chronic obstructive pulmonary disease	min	minute	SVT	supraventricular tachycardia
CPR	cardiopulmonary resuscitation	ml	milliliter	Sx	symptoms
CSF	cerebral spinal fluid	mm	millimeter	TIA	transient ischemic attack
CVA	cerebral vascular accident	MS	morphine sulfate	TID	three times a day
D/C	discontinue	NaCl	sodium chloride	TKO	to keep open
DOA	dead on arrival	NaHCO ₃	sodium bicarbonate	VF	ventricular fibrillation
D5W	5% dextrose in water	NG,N/G	nasogastric	w/s	watt second setting
Dx	diagnoses	nitro	nitroglycerine	x	times
ED	emergency department	NPO	nothing by mouth	y/o	years old
EKG/ECG	electrocardiogram	NS	normal saline		
Epi	epinephrine	NSR	normal sinus rhythm		
		NTG	nitroglycerine		
		O ₂	oxygen		
		OB	obstetrics		
		OD	overdose		
		OR	operating room		
		P	pulse		
		p	after		

Appendix M

Guidelines for New Protocol Development

*A rational decision making process**

Can also be used to evaluate existing protocols

Making a decision to develop a new protocol or evaluate an existing one should be based on a rational process. Questions that should be asked and answered when considering a new drug therapy or procedure are as follows:

Key Questions for any New Protocol

1. Is the drug therapy or procedure medically indicated and safe?
2. Is it within the scope of practice for the provider?
3. How specifically will this protocol benefit patient care?
4. What specifically is needed to implement this protocol (education/training, medical director protocol development/authorization, equipment needs, etc.)?
5. How will this protocol impact operation?
6. What is the opinion of providers concerning this protocol?
7. Does the medical community support this protocol change?
8. What are all the costs versus benefits associated with implementation and maintenance?
9. What are the medical-legal implications?
10. What ongoing provider involvement such as skills maintenance and continuous quality improvement is necessary?
11. How will success be measured?

Rational Protocol Development Process to Make the Right Protocol Decision

1. Study the issue thoroughly
2. Identify key questions
3. Compare with goals
4. Assess fit with system
5. Cost benefit analysis
6. Identify measuring tools

Stakeholders in this process are recognized to include, but not be limited to:

1. Medical direction (on-line and off-line)
2. Educators/training programs
3. Regulators of policy and rules
4. Service directors
5. Service providers
6. Consumers
7. Third party payers

*Developed based upon discussion at the October 1998 meeting of the Quality Assurance, Standards, and Protocols subcommittee of the Iowa EMS Advisory Council; and on concepts from the article 'When to Implement Clinical Protocol Change?' From EMS Best Practices September 1998.